

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 0 652 774 B1**

(12)

**EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention  
of the grant of the patent:  
**07.05.1997 Bulletin 1997/19**

(21) Application number: **93917118.7**

(22) Date of filing: **12.07.1993**

(51) Int. Cl.<sup>6</sup>: **A61K 47/32**, A61K 47/00

(86) International application number:  
**PCT/US93/06542**

(87) International publication number:  
**WO 94/02176 (03.02.1994 Gazette 1994/04)**

**(54) PHARMACEUTICAL COMPOSITION FOR TOPICAL USE CONTAINING A CROSSLINKED CATIONIC POLYMER AND AN ALKOXYLATED ETHER**

PHARMAZEUTISCHE ZUSAMMENSETZUNG ZUR TOPISCHEN ANWENDUNG DIE EIN VERNETZTES KATIONISCHES POLYMER UND EINEN ALKOXYLIERTE ÄTHER ENTHÄLT

COMPOSITION PHARMACEUTIQUE A EMPLOI TOPIQUE CONTENANT UN POLYMERE CATIONIQUE RETICULE ET UN ETHER ALCOXYLE

(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IE IT LI LU NL PT SE**

(30) Priority: **28.07.1992 US 920937**  
**25.09.1992 US 950527**  
**18.03.1993 US 33211**  
**25.06.1993 US 79977**

(43) Date of publication of application:  
**17.05.1995 Bulletin 1995/20**

(73) Proprietor: **THE PROCTER & GAMBLE COMPANY**  
**Cincinnati, Ohio 45202 (US)**

(72) Inventors:  
• **BLOOM, Roberta Christine**  
**Huntington, CT 06484 (US)**

• **DECKNER, George Endel**  
**Cincinnati, OH 45249 (US)**

(74) Representative: **Brooks, Maxim Courtney et al**  
**Procter & Gamble**  
**(Health & Beauty Care) Limited**  
**Rusham Park,**  
**Whitehall Lane**  
**Egham, Surrey TW20 9NW (GB)**

(56) References cited:  
**EP-A- 0 524 434** **WO-A- /00882**  
**WO-A-89/03669** **WO-A-93/07903**

• **PATENT ABSTRACTS OF JAPAN vol. 11, no. 361**  
**(C-459), 25 November 1987; & JP-A-62 132 812**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 0 652 774 B1**

## Description

The present invention relates to compositions for the topical administration of drugs, especially such compositions having enhanced penetration of the drug through the skin.

Because of the accessibility and large area of the skin, it has long been considered a promising route for the administration of drugs, whether dermal, regional, or systemic effects are desired.

The advantages of the topical route of drug administration include: avoidance of the risks and inconvenience of parenteral treatment; avoidance of the variable absorption and metabolism associated with oral treatment; continuity of drug administration, permitting use of pharmacologically active agents with short biological half-lives; potential reduction of gastrointestinal irritation in systemic administration; and treatment of cutaneous manifestations of diseases usually treated systemically.

However, the impermeability of skin is well-known, serving as a barrier to ingress of pathogens and toxic chemicals, and egress of physiologic fluids. This impermeability is the result of normal physiologic changes in developing skin. A typical cell in the epidermis is formed in the basal layer. It typically takes approximately thirty days for a cell to migrate from the basal layer of the epidermis to sloughing off and discarding at the outer layers of the stratum corneum. As the cell migrates outward from the basal layer, it progressively keratinizes until it is relatively impermeable. The result is the stratum corneum, an extremely thin surface layer (10 microns) with substantial barrier properties. The cell envelopes of the cells in the stratum corneum tend to be mainly polar lipids, such as ceramides, sterols, and fatty acids while the cytoplasm of stratum corneum cells remains polar and aqueous. Despite the close packing of the cells, some 15% of the stratum corneum is intercellular and, generally, lipid based. It is generally recognized that over the very short term, penetration occurs through the hair follicles and the sebaceous apparatus; long-term penetration occurs across cells (non-polar route). Poor penetration of many drugs across the epidermal lipid barrier has, until now, frustrated attempts to deliver clinically significant doses of many drugs by the topical route.

One route of delivery of drugs is by transdermal administration. Transdermal administration of drugs can be used in many instances to achieve therapeutic levels of the drugs in the systemic circulatory system, as well as for more localized internal dosing of drugs. Where such therapeutic levels of drugs can be achieved by transdermal administration, several potential advantages exist over other routes of administration. Sustained systemic delivery of drug controlled at therapeutic but below toxic levels over long periods of time with a single continuous application is often an advantage of transdermal drug administration. Potential contamination of internal tissues with undesired foreign substances or microbes, often associated with parenteral administration of drugs, is avoided with transdermal drug administration. Oral administration of many drugs is undesirable or unfeasible because the drug decomposes in the harsh environment of the gastrointestinal tract, lacks sufficient absorption from the gastrointestinal tract, or causes gastrointestinal upset or tissue damage in the gastrointestinal tract. First-pass metabolism of orally administered drugs can increase the dosage required to achieve therapeutic levels and thereby increase undesirable side effects either from the primary drug or the metabolites. Maintenance of uniform, optimal systemic levels of drugs for long periods of time is often difficult through oral administration. Such problems can often be reduced or avoided by transdermal drug administration.

Despite the substantial potential advantages for transdermal administration of drugs, relatively few drugs are so administered. The skin is a formidable barrier to the passage of most drugs. It is often necessary to provide a composition containing a skin penetration enhancing vehicle in order to provide sufficient transdermal penetration of the drug to achieve therapeutic levels of the drug at the target internal tissue. A number of skin penetration enhancing vehicles for drugs have been disclosed, including those in the following references: US-A-3,536,816 issued to Kellner on October 27, 1970; US-A-4,006,218 issued to Sipos on February 1, 1977; US-A-4,124,720 issued to Wenmaekers on November 7, 1978; US-A-4,126,681 issued to Reller on November 21, 1978; US-A-4,299,826 issued to Luedders on November 10, 1981; US-A-4,305,936 issued to Klein on December 15, 1981; US-A-4,309,414 issued to Inagi, Muramatsu & Nagai on January 5, 1982; US-A-4,338,306 issued to Kitao & Nishimura on July 6, 1982; US-A-4,442,090 issued to Kakeya, Kitao & Nishimura on April 10, 1984; US-A-4,485,033 issued to Kitao & Nishimura on November 27, 1984; US-A-4,537,776 issued to Cooper on August 27, 1985; US-A-4,552,872 issued to Cooper, Loomans & Fawzi on November 12, 1985; US-A-4,557,934 issued to Cooper on December 10, 1985; US-A-4,573,995 issued to Chen, Chun & Ensore on March 4, 1986; US-A-4,626,539 issued to Aungst & DiLuocio on December 2, 1986; US-A-4,637,930 issued to Konno, Kawata, Aruga, Sonobe & Mitomi issued January 20, 1987; US-A-4,695,465 issued to Kigasawa, Ohtani, Tanaka & Hayashida on September 22, 1987; EP-A-0,043,738 of The Procter & Gamble Company in the names of Wickett, Cooper & Loomans, published on June 13 1982; EP-A-0,095,813 of The Procter & Gamble Company in the name of Cooper, published December 7, 1983; PCT International Patent Application No. WO 87/03490 of Key Pharmaceuticals, Inc. in the names of Bodor and Loftson, published on June 18, 1987; Washitake, M., T. Anmo, I. Tanaka, T. Arita & M. Nakano, "Percutaneous Absorption of Drugs from Oily Vehicles", *Journal of Pharmaceutical Sciences*, Vol. 64, No. 3 (March, 1975), pp. 397-401; Shahi, V., & J. L. Zatz, "Effect of Formulation Factors on Penetration of Hydrocortisone through Mouse Skin", *Journal of Pharmaceutical Sciences*, Vol. 67, No. 6 (June, 1978), pp. 789-792; Cooper, E.R., "Increased Skin Permeability for Lipophilic Molecules", *Journal of Pharmaceutical Sciences*, Vol. 73, No. 8

(August, 1984), pp. 1153-1156; Aungst, B.J., N. J. Rogers & E. Shefter, "Enhancement of Naloxone Penetration through Human Skin *In Vitro* Using Fatty Acids, Fatty Alcohols, Surfactants, Sulfoxides and Amides", *International Journal of Pharmaceutics*, Vol. 33 (1986), pp. 225-234; Green, P.G., & J. Hadgraft, "Facilitated Transfer of Cationic Drugs Across a Lipoidal Membrane by Oleic Acid and Lauric Acid", *International Journal of Pharmaceutics*, Vol. 37 (July, 1987), pp. 251-255.

Often, though, transdermal delivery of drugs can cause skin irritation (e.g., burning and stinging), drying of skin and further can be difficult to formulate. The present invention provides stable compositions for the transdermal delivery of drugs which do not require ethanol to solubilize the active and therefore can be substantially free of ethanol, are easy to formulate and which further provide low dermal irritation. These compositions also provide improved moisturizing benefits.

It is an object of the present invention to provide novel compositions for enhancing the skin penetration of drugs with significantly reduced skin irritation.

It is a further object of the present invention to provide such compositions which provide sufficient skin penetration enhancement to achieve therapeutic levels of the drugs in target internal tissues.

It is a further object of the present invention to provide such compositions with low dermal irritation, especially in compositions requiring a low pH.

It is a further object of the present invention to provide such compositions having good stability and good cosmetics and which do not require ethanol to solubilize the drug actives.

It is a further object of the present invention to provide compositions useful for the treatment of acne.

It is a further object of the present invention to provide compositions useful for the treatment of skin wrinkles and skin atrophy and other affects of skin aging such as actinic elastosis.


According to the present invention there is provided a pharmaceutical composition for topical application having enhanced penetration through the skin comprising:

(a) a safe and effective amount of a pharmaceutical active;

(b) from 0.1% to 10.0% of a high molecular weight crosslinked cationic polymer of the formula:  $(A)_l(B)_m(C)_n$  wherein (A) is a dialkylaminoalkyl acrylate monomer or its quaternary ammonium or acid addition salt, (B) is a dialkylaminoalkyl methacrylate monomer or its quaternary ammonium or acid addition salt, (C) is a monomer having one carbon-carbon double bond, l is an integer of 0 or greater, m is an integer of 1 or greater, and n is an integer of 0 or greater, wherein said polymer contains a crosslinking agent;

(c) from 0.05% to 5% of a high HLB non-ionic surfactant selected from ethoxylated alcohols having from 8 to 22 carbons and from 6 to 30 ethylene oxide groups and

(d) from 0.1% to 25% of an alkoxylated ether of the formula:



wherein R is straight or branched chain  $C_1$  to  $C_{18}$ , preferably  $C_1$  to  $C_{10}$  and most preferably  $C_1$  to  $C_5$ , and X is methyl or ethyl, and wherein n is an average of from 5 to 20; wherein the composition is substantially free of ethanol.

In further embodiments the compositions comprise from 0.01% to 5% of a low HLB non-ionic surfactant.

In even further embodiments the crosslinked cationic polymer is of the formula wherein (C) is acrylamide.

In even further embodiments the crosslinked cationic polymer is of the formula wherein (C) is acrylamide and l is zero.

In even further embodiments the crosslinked cationic polymer is a homopolymer wherein both l and n are zero.

All concentrations and ratios herein are by weight of total composition and all measurements are at 25°C, unless otherwise specified.

These compositions are substantially free of ethanol. By substantially free of ethanol is meant the compositions comprise less than about 5%, preferably less than about 1% and most preferably less than about 0.5% ethanol.

The invention hereof can comprise, consist of, or consist essentially of, the essential as well as optional ingredients and components described herein.

The present invention involves compositions comprising certain specific cationic polymers, nonionic surfactant and water-immiscible solubilizing aids which may be applied topically to the skin and which result in improved transdermal delivery (i.e. optimized penetration) of the drugs through the skin. These compositions do not need additional solvents such as ethanol to solubilize the active in the compositions.

**Drug Active** The compositions of the present invention comprise a safe and effective amount of a drug active. The phrase "safe and effective amount", as used herein, means an amount of a drug high enough to significantly positively

modify the condition to be treated, but low enough to avoid serious side effects (at a reasonable benefit/risk ratio), within the scope of sound medical judgement. A safe and effective amount of the drug will vary with the specific drug, the ability of the composition to penetrate the drug through the skin, the amount of composition to be applied, the particular condition being treated, the age and physical condition of the patient being treated, the severity of the condition, the duration of the treatment, the nature of concurrent therapy, and like factors.

The drug compounds present in the compositions of the present invention preferably comprise from about 0.1% to about 20% by weight of the compositions, more preferably from about 0.1% to about 10%, and most preferably from about 0.1% to about 5%. Mixtures of drug actives may also be used. It is contemplated herein that the various actives described below can provide more than one benefit and can alternatively be classified under more than one category, below.

Useful drug actives in the compositions of the present invention include anti-acne drugs. Anti-acne drugs preferred for use in the present invention include the keratolytics such as salicylic acid, sulfur, lactic acid, glycolic, pyruvic acid, urea, resorcinol, and N-acetylcysteine; retinoids such as retinoic acid and its derivatives (e.g., cis and trans); antibiotics and antimicrobials such as benzoyl peroxide, octopirox, erythromycin, zinc, tetracyclin, triclosan, azelaic acid and its derivatives, phenoxy ethanol and phenoxy propanol, ethylacetate, clindamycin and meclocycline; sebastats such as flavinoids; alpha and beta hydroxy acids; and bile salts such as scymnol sulfate and its derivatives, deoxycholate, and cholate. Preferred for use herein is salicylic acid. Without being limited by theory, even though salicylic acid is classified herein as a an anti-acne agent, it is also useful for other skin benefits such as anti-wrinkling and anti-skin atrophy benefits which are described in the following paragraph.

Useful drug actives in the compositions of the present invention include anti-wrinkle and anti-skin atrophy actives which include the C2-C30 alpha-hydroxy acids (e.g., glycolic acid, lactic acid, 2-hydroxybutanoic acid, and the like), retinoic acid, salicylic acid, and skin peel agents (e.g., phenol, acetic acid, and the like). Preferred for use herein are glycolic acid, lactic acid, salicylic acid, and mixtures thereof.

Useful drug actives in the compositions of the present invention include non-steroidal anti-inflammatory drugs (NSAIDS). The NSAIDS can be selected from the following categories: propionic acid derivatives; acetic acid derivatives; fenamic acid derivatives; biphenylcarboxylic acid derivatives; and oxicams. All of these NSAIDS are fully described in the US-A-4,985,459 to Sunshine et al., issued January 15, 1991. Most preferred are the propionic NSAIDS including but not limited to aspirin, acetaminophen, ibuprofen, naproxen, benoxaprofen, flurbiprofen, fenoprofen, fenbuten, ketoprofen, indoprofen, pirofen, carprofen, oxaprozin, pranoprofen, miroprofen, tiroxaprofen, suprofen, alminoprofen, tiaprofenic acid, fluprofen and bucloxic acid. Also useful are the steroidal anti-inflammatory drugs including hydrocortisone and the like.

Useful drug actives in the compositions of the present invention include antihistaminic drugs. Antihistaminic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of chlorpheniramine, triprolidine, diphenhydramine, doxylamine, pyrilamine, phenindamine, promethazine, cyproheptadine, azatadine, clemastine, carbinoxamine, tripeleminamine, terfenadine, dexchlorpheniramine, brompheniramine, chlorcyclizine, diphenylpyraline, pheniramine and phenyltoloxamine, and mixtures thereof.

Useful drug actives in the compositions of the present invention include antitussive drugs. Antitussive drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of dextromethorphan, codeine, caramiphen and carbetapentane.

Useful drug actives in the compositions of the present invention include antipruritic drugs. Antipruritic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of methdilazine and trimeprazine.

Useful drug actives in the compositions of the present invention include anticholinergic drugs. Anticholinergic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of scopolamine, atropine, homatropine, levodopa, dicyclomine, hyoscyamine, procyclidine, trihexyphenidyl and ethopropazine.

Useful drug actives in the compositions of the present invention include anti-emetic and antinauseant drugs. Anti-emetic and antinauseant drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of cyclizine, medizine, chlorpromazine, buclizine, metoclopramide, prochlorperazine and trimethobenzamide.

Useful drug actives in the compositions of the present invention include anorexic drugs. Anorexic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of benzphetamine, phentermine, chlorphentermine, fenfluramine, diethylpropion and phendimetrazine.

Useful drug actives in the compositions of the present invention include central stimulant drugs. Central stimulant drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of amphetamine, methamphetamine, dextroamphetamine and methylphenidate.

Useful drug actives in the compositions of the present invention include antiarrhythmic drugs. Antiarrhythmic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of propranolol, procainamide, disopyramide, quinidine, encainide, flecanaide, mexiletine and tocainide. Other antiarrhythmic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of the

quinidine derivatives disclosed in US-A-4,716,171 issued to Jarreau and Koenig on December 29, 1987. Highly preferred compounds included in this class include pharmaceutically-acceptable salts of 3S-hydroxy-10,11-dihydroquinidine, 3R-hydroxy-10,11-dihydroquinidine, 3R-hydroxy-0-acetyl-10,11-dihydroquinidine, and 3S-hydroxy-0-acetyl-10,11-dihydroquinidine, especially 3S-hydroxy-10,11-dihydroquinidine.

5 Useful drug actives in the compositions of the present invention include  $\beta$ -adrenergic blocker drugs.  $\beta$ -Adrenergic blocker drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of metoprolol, acebutolol, betaxolol, labetalol and timolol.  $\beta$ -Adrenergic blocker drugs more preferred for inclusion in compositions of the present invention include metoprolol tartrate, acebutolol hydrochloride, betaxolol hydrochloride, labetalol hydrochloride and timolol maleate.

10 Useful drug actives in the compositions of the present invention include cardiotonic drugs. Cardiotonic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of milrinone, amrinone and dobutamine. Other cardiotonic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of 14-amino steroid derivatives, some of which are disclosed in U.S. Patent Nos. 4,325,879, 4,552,868 and 4,584,289, issued to Jarreau and Koenig on April 20, 1982, November 12, 1985 and April 22, 1986, respectively.

Useful drug actives in the compositions of the present invention include antihypertensive drugs. Antihypertensive drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of enalapril, clonidine, hydralazine, minoxidil (which is also a hair growth stimulator drug), guanadrel, guanethidine, guanfacine, mecamlamine, methyldopate, pargyline, phenoxylbenzamine and prazosin.

20 Useful drug actives in the compositions of the present invention include diuretic drugs. Diuretic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of amiloride and hydrochlorothiazide. Diuretic drugs more preferred for inclusion in compositions of the present invention include amiloride hydrochloride.

Useful drug actives in the compositions of the present invention include vasodilator drugs. Vasodilator drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of diltazem, amiodarone, isoxsuprine, nylidrin, tolazoline and verapamil.

Useful drug actives in the compositions of the present invention include vasoconstrictor drugs. Vasoconstrictor drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of dihydroergotamine, ergotamine and methysergide.

30 Useful drug actives in the compositions of the present invention include anti-ulcer drugs. Anti-ulcer drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of ranitidine and cimetidine.

Useful drug actives in the compositions of the present invention include anesthetic drugs. Anesthetic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of lidocaine, bupivacaine, chlorprocaine, dibucaine, etidocaine, mepivacaine, tetracaine, dyclonine, hexylcaine, procaine, cocaine, ketamine, pramoxine and phenol.

Useful drug actives in the compositions of the present invention include antidepressant drugs. Antidepressant drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of imipramine, desipramine, amitriptyline, nortriptyline, protriptyline, doxepin, maprotiline, phenelzine, tranylcypromine, trazodone and trimipramine.

Useful drug actives in the compositions of the present invention include tranquilizer and sedative drugs. Tranquilizer and sedative drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of chlordiazepoxide, benactyzine, benzquinamide, flurazepam, hydroxyzine, loxapine and promazine.

Useful drug actives in the compositions of the present invention include anti psychotic drugs. Antipsychotic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of chlorpromazine, fluphenazine, haloperidol, molindone, thioridazine and trifluoperazine.

Useful drug actives in the compositions of the present invention include antimicrobial drugs (antibacterial, antifungal, antiprotozoal and antiviral drugs). Antimicrobial drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of  $\beta$ -lactam drugs, quinolone drugs, ciprofloxacin, norfloxacin, tetracycline, erythromycin, amikacin, triclosan, doxycycline, capreomycin, chlorhexidine, chlortetracycline, oxytetracycline, clindamycin, ethambutol, hexamidine isethionate, metronidazole, pentamidine, gentamicin, kanamycin, lineomycin, methacycline, methenamine, minocycline, neomycin, netilmicin, paromomycin, streptomycin, tobramycin, miconazole and amantadine. Antimicrobial drugs preferred for inclusion in compositions of the present invention include tetracycline hydrochloride, erythromycin estolate, erythromycin stearate (salt), amikacin sulfate, doxycycline hydrochloride, capreomycin sulfate, chlorhexidine gluconate, chlorhexidine hydrochloride, chlortetracycline hydrochloride, oxytetracycline hydrochloride, clindamycin hydrochloride, ethambutol hydrochloride, metronidazole hydrochloride, pentamidine hydrochloride, gentamicin sulfate, kanamycin sulfate, lineomycin hydrochloride, methacycline hydrochloride, methenamine hippurate, methenamine mandelate, minocycline hydrochloride, neomycin sulfate, netilmicin sulfate, paromomycin sulfate, streptomycin sulfate, tobramycin sulfate, miconazole hydrochloride, amantadine hydrochloride ,

amanafadine sulfate, triclosan, octopirox, parachlorometa xylenol, nystatin, tolnaftate and clotrimazole.

Useful drug actives in the compositions of the present invention include antineoplastic drugs. Antineoplastic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of bleomycin, daunorubicin, doxorubicin, mechlorethamine, procarbazine, quinacrine, tamoxifen, vinblastine and vincristine.

5 Useful drug actives in the compositions of the present invention include antimalarial drugs. Antimalarial drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of chloroquine, hydroxychloroquine primaquine and quinine.

Useful drug actives in the compositions of the present invention include muscle relaxant drugs. Muscle relaxant drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of cinnamedrine, cyclobenzaprine, flavoxate, orphenadrine, papaverine, mebeverine, idaverine, ritodrine, dephenoxylate, dantrolene and azumolene.

Useful drug actives in the compositions of the present invention include antispasmodic drugs. Antispasmodic drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of the compounds disclosed in US-A-3,856,825 issued to Wright, Burch and Goldenburg on December 24, 1974.

15 Useful drug actives in the compositions of the present invention include antidiarrheal drugs. Antidiarrheal drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of loperamide.

Useful drug actives in the compositions of the present invention include bone-active drugs. Bone-active drugs preferred for inclusion in compositions of the present invention include pharmaceutically-acceptable salts of diphosphonate drug compounds and phosphonoalkylphosphinate drug compounds, including the prodrug esters thereof. Such compounds are disclosed, for example, in US-A-3,683,080 issued to Francis on August 8, 1972; US-A-4,304,734 issued to Jary, Rihakova & Zobacova on December 8, 1981; US-A-4,687,768 issued to Benedict & Johnson on August 18, 1987; US-A-4,711,880 issued to Stahl & Schmitz on December 8, 1987; and US-A-4,719,203 issued to Bosies & Gall on January 12, 1988; copending U.S. patent application Serial Nos. 808,584, of Benedict & Perkins filed December 13, 1985; 25 945,069 of Ebetino, Buckingham & McOske filed December 19, 1986; 945,068 of Ebetino & Benedict filed December 19, 1986; and 069,666 of Ebetino filed July 6, 1987; and EP-A-0,001,584 of Blum, Hempel & Worms, published May 2, 1979; EP-A-0,039,033 published April 11, 1981; EP-A-0,186,405 of Benedict & Perkins, published July 2, 1986; and EP-A-0,243,173 of Oku, Todo, Kasahara, Nakamura, Kayakiri & Hashimoto, published October 28, 1987.

Also useful in the present invention sunscreens agents. A wide variety of sunscreens agents are described in 30 US-A-5,087,445, to Haffey et al., issued February 11, 1992; US-A-5,073,372, to Turner et al., issued December 17, 1991; US-A-5,073,371, to Turner et al. issued December 17, 1991; and Segarin, et al., at Chapter VIII, pages 189 et seq., of Cosmetics Science and Technology.

Preferred among those sunscreens which are useful in the compositions of the instant invention are those selected from the group consisting of 2-ethylhexyl p-methoxycinnamate, 2-ethylhexyl N,N-dimethyl-p-aminobenzoate, p-aminobenzoic acid, 2-phenylbenzimidazole-5-sulfonic acid, octocrylene, oxybenzone, homomenthyl salicylate, octyl salicylate, 4,4'-methoxy-1-butylidibenzoylmethane, 4-isopropyl dibenzoylmethane, 3-benzylidene camphor, 3-(4-methylbenzylidene) camphor, titanium dioxide, zinc oxide, silica, iron oxide, and mixtures thereof.

Still other useful sunscreens are those disclosed in US-A-4,937,370, to Sabatelli, issued June 26, 1990; and US-A-4,999,186, to Sabatelli et al., issued March 12, 1991. The sunscreens agents disclosed therein have, in a single 40 molecule, two distinct chromophore moieties which exhibit different ultra-violet radiation absorption spectra. One of the chromophore moieties absorbs predominantly in the UVB radiation range and the other absorbs strongly in the UVA radiation range. These sunscreens agents provide higher efficacy, broader UV absorption, lower skin penetration and longer lasting efficacy relative to conventional sunscreens. Especially preferred examples of these sunscreens include those selected from 4-N,N-(2-ethylhexyl)methylaminobenzoic acid ester of 2,4-dihydroxybenzophenone, 4-N,N-(2-ethylhexyl)methylaminobenzoic acid ester with 4-hydroxydibenzoylmethane, 4-N,N-(2-ethylhexyl)methylaminobenzoic acid ester of 2-hydroxy-4-(2-hydroxyethoxy)benzophenone, 4-N,N-(2-ethylhexyl)-methylaminobenzoic acid ester of 4-(2-hydroxyethoxy)dibenzoylmethane, and mixtures thereof.

Generally, the sunscreens can comprise from about 0.5% to about 20% of the compositions useful herein. Exact amounts will vary depending upon the sunscreen chosen and the desired Sun Protection Factor (SPF). SPF is a commonly used measure of photoprotection of a sunscreen against erythema. See Federal Register, Vol. 43, No. 166, pp. 38206-38269, August 25, 1978.

Also useful in the present invention are sunless tanning agents including dihydroxyacetone, glyceraldehyde, indoles and their derivatives, and the like. These sunless tanning agents can also be used in combination with the sunscreen agents.

55 Other useful actives include skin bleaching (or lightening) agents including but not limited to hydroquinone, ascorbic acid, kojic acid and sodium metabisulfite and wound healing agents such as peptide derivatives, yeast, panthenol, lamin and kinetin.

Water-Soluble Polymer The polymers useful in the present invention are certain cationic polymers. These polymers are generally described in US-A-5,100,660, to Haw et al., issued March 31, 1992; US-A-4,849,484, to Heard, issued

July 18, 1989; US-A-4,835,206, to Farrar et al., issued May 30, 1989; US-A-4,628,078 to Glover et al. issued December 9, 1986; US-A-4,599,379 to Flesher et al. issued July 8, 1986; and EP-A-0,228,868 to Farrar et al., published July 15, 1987.

The compositions of the instant invention comprise from 0.1% to 10%, preferably from 0.1% to 7.5%, and most preferably from 0.1% to 5% of the polymer.

In general these polymers are high molecular weight materials containing cationic, usually quaternized, nitrogen moieties. These polymers can be characterized by the general formula:  $(A)_l(B)_m(C)_n$  wherein (A) is a dialkylaminoalkyl acrylate monomer or its quaternary ammonium or acid addition salt, (B) is a dialkylaminoalkyl methacrylate monomer or its quaternary ammonium or acid addition salt, (C) is a monomer having one carbon-carbon double bond, l is an integer of 0 or greater, m is an integer of 1 or greater, and n is an integer of 0 or greater. The (C) monomer can be selected from any of the commonly used monomers. Nonlimiting examples of these monomers include ethylene, propylene, butylene, isobutylene, eicosene, maleic anhydride, acrylamide, methacrylamide, maleic acid, acrolein, cyclohexene, ethyl vinyl ether, and methyl vinyl ether. In the cationic polymers of the present invention, (C) is preferably acrylamide.

In highly preferred embodiments, these polymers also contain a crosslinking agent, which is most typically a material containing one or more unsaturated functional groups. In other words, these crosslinked polymers are expressly intended to also include the crosslinking agent in addition to the (A), (B), and (C) monomer units. Nonlimiting examples of suitable crosslinking agents include those selected from methylenebisacrylamides, diallyldialkyl ammonium halides, polyalkenyl polyethers of polyhydric alcohols, allyl acrylates, vinyloxyalkylacrylates, and polyfunctional vinylidenes. Specific examples of crosslinking agents useful herein include those selected from methylenebisacrylamide, ethylene glycol di-(meth)acrylate, di-(meth)acrylamide, cyanomethylacrylate, vinyloxyethylmethacrylate, allyl pentaerythritol, trimethylolpropane diallylether, allyl sucrose, butadiene, isoprene, divinyl benzene, divinyl naphthalene, ethyl vinyl ether, methyl vinyl ether, and allyl acrylate. Other crosslinkers include formaldehyde and glyoxal. Preferred for use herein as a crosslinking agent is methylenebisacrylamide.

When the crosslinking agent is present, widely varying amounts can be employed depending upon the properties desired in the final polymer, e.g. viscosifying effect. Without being limited by theory, it is believed that incorporation of a crosslinking agent into these cationic polymers provides a material that is a more effective viscosifying agent without negatives such as stringiness and viscosity breakdown in the presence of electrolytes. The crosslinking agent, when present, can comprise from about 1 ppm to about 1000 ppm, preferably from about 5 ppm to about 750 ppm, more preferably from about 25 ppm to about 500 ppm, even more preferably from about 100 ppm to about 500 ppm, and most preferably from about 250 ppm to about 500 ppm of the total weight of the polymer on a weight/weight basis.

In one group of embodiments, these cationic polymers are made from processes which generally require polymerisation of a solution containing from about 20% to about 60%, generally from about 25% to about 40%, by weight monomer, in the presence of an initiator (usually redox or thermal) until the polymerization terminates. The temperature generally starts low, e.g. 0° to 95°C. The polymerization can be conducted by forming a reverse phase dispersion of an aqueous phase of the monomers into a nonaqueous liquid, e.g. mineral oil and the like.

When the polymer contains acrylamide, the molar proportion of acrylamide, based on the total molar amount of acrylamide, dialkylaminoalkyl acrylate and dialkylaminoalkyl methacrylate, is generally from about 20% to about 99%. Preferably, the amount of acrylamide is at least 50%, often at least 60% to below about 95%. All percentages describing the polymer herein are molar, unless otherwise specified.

Where monomer A is present, the ratio of monomer A:monomer B used in this process, and thus the ratio of groups A and B in the final polymer, on a molar basis is preferably 80:20 to 20:80. In one class of processes, the ratio is 5:95 to 50:50, i.e., the cationic monomer is mainly methacrylate. In these processes, the ratio is generally being achieved in the range of from 25:75 to 5:95.

In another class of processes, the ratio A:B is from 50:50 to 85:15, the cationic monomers being mainly acrylate. Preferably the ratio A:B is 60:40 to 85:15, most preferably 75:25 to 85:15.

Preferred is where monomer A is not present and the ratio of monomer B:monomer C is from 30:70 to 70:30, preferably from 40:60 to 60:40 and most preferably from 45:55 to 55:45.

The polymerisation is preferably conducted under known conditions such that the polymers are water soluble and have a high molecular weight, generally about 1 million, for instance up to 30 million. The intrinsic viscosity, measured in molar sodium chloride solution at 25° C., is generally above 6, for instance from 8 to 14.

A cationic polymer useful herein is one conforming to the general structure  $(A)_l(B)_m(C)_n$  wherein l is zero, (B) is methyl quaternized dimethylaminoethyl methacrylate, the ratio of (B):(C) is 45:55 to 55:45, and the optional crosslinking agent is methylenebisacrylamide. This polymer which has the proposed CTFA designation, "Polyquaternium 32 (and Mineral Oil)", is commercially available as Salcare<sup>(RTM)</sup> SC92 from Allied Colloids Ltd. (Norfolk, VA).

Alternatively in another group of preferred embodiments, these cationic polymers do not contain the acrylamide monomer, that is, n is zero. In these polymers the (A) and (B) monomer components are as described above. An especially preferred group of these non-acrylamide containing polymers is one in which l is also zero. In this instance the polymer is essentially a homopolymer of a dialkylaminoalkyl methacrylate monomer or its quaternary ammonium or acid addition salt. These dialkylaminoalkyl methacrylate copolymers and homopolymers preferably contain a crosslink-



ing agent as described above.

A cationic homopolymer useful herein is one conforming to the general structure  $(A)_l(B)_m(C)_n$  wherein  $l$  is zero,  $(B)$  is methyl quaternized dimethylaminoethyl methacrylate,  $n$  is zero, and the crosslinking agent is methylenebisacrylamide. This polymer, which has recently been given CTFA designation "Polyquaternium 37 (and) Mineral Oil (and) PPG-1 Trideceth-6", will be referred to herein as crosslinked methyl quaternized dimethylaminoethyl methacrylate and mineral oil. This polymer is commercially available as Salcare<sup>(RTM)</sup> SC95 from Allied Colloids Ltd. (Norfolk, VA).

#### Nonionic Surfactant

Compositions of this invention also contain from 0.05% to 10%, preferably from 0.05% to 5%, more preferably from 0.05% to 2%, and most preferably from 0.25% to 1% of certain nonionic surfactants. Nonionic surfactants used herein have an HLB of from about 6 to about 18, preferably from about 8 to about 18, and more preferably from about 10 to about 18. These nonionic surfactants are herein designated as "high HLB nonionic surfactants" and do not include the surfactants with HLB values less than 6, as described below.

The abbreviation "HLB" stands for hydrophilic lipophilic balance. The HLB system is well known in the art and is described in detail in "The HLB System, A Time-Saving Guide to Emulsifier Selection", ICI Americas Inc., August 1984.

High HLB nonionic surfactants used herein are ethoxylated alcohols having a straight or branched chain alkyl group in the  $C_{8-22}$ , preferably  $C_{10-20}$ , more preferably  $C_{12-20}$ , range and from about 6 to about 30, preferably from about 6 to about 25, ethylene oxide groups.

Preferred for use herein are polyethylene oxide ethers derived from lauryl alcohol, cetyl alcohol, oleyl alcohol, stearyl alcohol, isostearyl alcohol, myristyl alcohol, behenyl alcohol, and mixtures thereof. More preferred for use herein, are: polyoxyethylene 10 cetyl ether, known by the CTFA designation as ceteth-10; polyoxyethylene (21) stearyl ether, known by the CTFA designation steareth-21; coconut alkyl polyethoxylate (6.5); decyl polyethoxylate (6); and mixtures thereof. Most preferred for use herein are ceteth-10, steareth-21, and mixtures thereof.

A detailed listing of suitable nonionic surfactants, of the above types, for the compositions herein can be found in US-A-4,557,853, to Collins, issued Dec. 10, 1985. Commercial sources of such surfactants can be found in McCutcheon's EMULSIFIERS AND DETERGENTS, North American Edition, 1984, McCutcheon Division, MC Publishing Company.

Especially preferred for use herein are compositions which contain at least one of the above described high HLB nonionic surfactants, in combination with at least one other nonionic surfactant having an HLB from about 1 to about, but not greater than or equal to 6, hereinafter designated as a "low HLB nonionic surfactant". These low HLB nonionic surfactants do not include the high HLB nonionic surfactants described above. Without being limited by theory, it is believed that this combination of both high and low HLB nonionic surfactants provides compositions demonstrating enhanced emulsion stability.

In these compositions utilizing both the low HLB nonionic surfactant in combination with the high HLB nonionic surfactant, the low HLB nonionic surfactant comprises from 0.01% to 5%, preferably from 0.01% to 1%, and more preferably from 0.025% to 0.5% of the composition.

Typical of these low HLB nonionic surfactants are ethoxylated alcohols with, the alcohol derivatives being preferred. In general, these alcohol derivatives contain a straight or branched chain alkyl group in the  $C_{8-22}$ , preferably  $C_{10-20}$ , more preferably  $C_{12-20}$ , range, and generally contain from 1 to 5 ethylene oxide groups per molecule.


Some nonlimiting examples of these low HLB nonionic surfactants useful herein include stearic acid ethoxylated with 1 mole of ethylene oxide (i.e. steareth-1), steareth-2, steareth-3, steareth-4, steareth-5, ceteth-1, cetheth-2, ceteth-3, ceteth-4, ceteth-5, laureth-1, laureth-2, laureth-3, laureth-4, laureth-5, oleic acid ethoxylated with 1 mole of ethylene oxide (i.e. oleth-1), oleth-2, oleth-3, oleth-4, oleth-5, and mixtures thereof. Preferred low HLB nonionic surfactants are steareth-1, steareth-2, steareth-3, ceteth-1, ceteth-2, cetheth-3, laureth-1, laureth-2, laureth-3, oleth-1, oleth-2, oleth-3, and mixtures thereof. More preferred are steareth-2, ceteth-2, laureth-2, oleth-2, and mixtures thereof. Most preferred is steareth-2, which is available commercially as Brij<sup>(RTM)</sup> 72 from ICI Americas.

Detailed listings of low HLB surfactants can be found in McCutcheon's EMULSIFIERS AND DETERGENTS, North American Edition, 1984, McCutcheon Division, MC Publishing Company.

When a low HLB nonionic surfactant is used herein in combination with a high HLB surfactant, it has been found that the combination of steareth-2 with ceteth-10 and/or steareth-21 is especially preferred.

Water-Immiscible Solubilizing Aid The compositions of the present invention comprise from 0.1% to 25%, more preferably from 0.1% to 15%, and most preferably from 6% to 10% of a water-immiscible solubilizing aid for the active ingredients useful herein. By water-immiscible is meant that the material has a solubility in water of less than 1 gram per 100 grams of water at 25°C. Without being limited by theory, these materials improve the stability of the composition as well as provide moisturization benefits. The water-immiscible solubilizing aids used herein is an alkoxyated ether. The alkoxyated ethers useful herein are of the formula:





wherein R is straight or branched chain C<sub>1</sub> to C<sub>18</sub>, preferably C<sub>1</sub> to C<sub>5</sub>, and most preferably C<sub>1</sub> to C<sub>4</sub>, and X is methyl or ethyl and n has an average value of from 5 to 20, preferably from 8 to 16, and more preferably from 10 to 16. Also preferred for use herein are the polyoxypropylene butyl ethers (i.e. where R is C<sub>4</sub>). These compounds are also known as PPG butyl ethers. A preferred compound among these butyl ether derivatives is PPG-14 butyl ether (i.e. a butyl ether having about 14 propylene oxide units incorporated in its structure). PPG-14 butyl ether is available as Fluid AP from Union Carbide. Alternatively, another preferred compound is PPG-15 stearyl ether (i.e. a stearyl ether having about 15 propylene oxide units incorporated in its structure). PPG-15 stearyl ether is available as Arlamol<sup>(TM)</sup> S3 and S7 from ICI Americas.

The polyalkylene oxide polymers or copolymers useful herein include polypropylene oxides, polybutylene oxides, mixed polyethylene and polypropylene oxides, mixed polyethylene and polybutylene oxides, and the like, provided that the water-immiscibility requirement is met. Nonlimiting examples of these polymers include PPG-9, PPG-15, PPG-17, PPG-20, PPG-26, PPG-30, PPG-34, a copolymer of ethylene oxide and propylene oxide wherein the copolymer has an average of 3 ethylene oxide units and an average of 20 propylene oxide units, and a copolymer of propylene oxide and butylene oxide wherein the copolymer has an average of 10 propylene oxide units and an average of 5 butylene oxide units.

In other embodiments, the water-immiscible solubilizing aids include those of the above chemical formula wherein R is a straight or branched chain C<sub>1</sub>-C<sub>8</sub> moiety having one or more free hydroxyl groups, wherein this moiety can be derived from the corresponding alcohol. Nonlimiting examples of these C<sub>1</sub>-C<sub>8</sub> moieties include those derived from glycerin, propylene glycol, butylene glycol, hexylene glycol, 1,2,6-hexanetriol, and the like.

**Vehicle** The compositions of the present invention are used along with pharmaceutically-acceptable carrier (or vehicle) components. The term "pharmaceutically-acceptable carrier components", as used herein, means compatible solid or liquid filler diluents which are suitable for administration to a human or lower animal. The term "compatible", as used herein, means that the components are capable of being commingled with the drug compounds, and other components of the compositions of the present invention, in a manner such that there is no interaction which would substantially reduce the pharmaceutical efficacy of the compositions of the present invention under ordinary use situations.

Pharmaceutically-acceptable carrier components must, of course, be of sufficiently high purity and sufficiently low toxicity to render them suitable for administration to the human or lower animal being treated.

Some examples of substances which can serve as pharmaceutically-acceptable carrier components are glycerol; ethanol; water; antioxidants; surfactants; chelating agents; preservatives; thickeners; anti-bacterial agents; as well as other non-toxic compatible substances used in pharmaceutical formulations.

These compositions can also contain one or more additional humectants/moisturizers, many of which may also be useful as actives. A variety of humectants/moisturizers can be employed and can be present at a level of from 0.5% to 30%, more preferably from 2% to 8%, and most preferably from 3% to 5%. These materials include polyhydroxy alcohols such as sorbitol, glycerin, hexanetriol, hexylene glycol and the like; sugars and starches; sugar and starch derivatives (e.g. alkoxylated glucose); D-panthenol and its derivatives; hyaluronic acid; lactamide monoethanolamine; acetamide monoethanolamine; and mixtures thereof. Other useful humectants include propylene oxide ethers of glycerin, preferably those having less than about four moles of propylene oxide per mole of glycerin. Useful examples of these propylene oxide ethers of glycerin are described in detail in US-A-4,976,953, to Orr et al., issued December 11, 1990.

Preferred humectants/moisturizers for use in the compositions of the present invention are the C<sub>3</sub>-C<sub>6</sub> diols and triols. Especially preferred is the triol, glycerin.

The compositions of the present invention can also optionally comprise at least one emollient. Examples of suitable emollients include, but are not limited to, volatile and non-volatile silicone oils, highly branched hydrocarbons, and mixtures thereof. Emollients useful in the instant invention are further described in US-A-4,919,934, to Deckner et al., issued April 24 1990.

The emollients can typically comprise in total from 1% to 50%, preferably from 1% to 25%, and more preferably from 1% to 10% by weight of the compositions of the present invention.

The compositions of this invention may also contain pharmaceutically acceptable optional components that modify the physical and/or therapeutic effects of the compositions. Such optional components may include, for example, additional solvents, gelling agents, fragrances, preservatives, anti-bacterial agents, thickeners (e.g., fatty alcohols such as cetyl alcohol and stearyl alcohol), and stabilizers. However, such optional materials must not unduly interfere with the transdermal delivery of the drug active. Optional components useful in the compositions of this invention are described in the following patent documents: EP-A-0,043,738, Wickett et al., published January 13, 1982; and US-A-4,552,872, Cooper et al., issued November 12, 1985.

Most preferred compositions herein are creams and lotions.

Another optional material is a solvent or co-solvent material.

Most preferred compositions herein have a pH of below about 5, preferably below about 4, and most preferably below about 3. Without being limited by theory, the pH of a formulation can be an important factor in the delivery and availability of an active ingredient. For example, for the active ingredient salicylic acid, at pH values above its  $pK_a$  in a particular matrix, the salicylic acid would exist primarily in its ionized form and would not as readily penetrate into the skin. Thus, an acidic formulation range is preferred for salicylic acid compositions in order to suppress ionization and enhance its penetration into the stratum corneum.

A wide variety of acids, bases, buffers, and sequestrants can be utilized to adjust and/or maintain the pH and ionic strength of the compositions useful in the instant invention. Materials useful for adjusting and/or maintaining the pH and/or the ionic strength include sodium carbonate, sodium hydroxide, hydrochloric acid, phosphoric acid, sulfuric acid, acetic acid, sodium acetate, sodium hydrogen phosphate, sodium dihydrogen phosphate, citric acid, sodium citrate, sodium bicarbonate, triethanolamine, EDTA, disodium EDTA, tetrasodium EDTA, and the like.

## Test Method

Transdermal penetration of drugs is conveniently determined and compared from various vehicles using the apparatus and procedure described below.

Full thickness excised human thigh skin is obtained from cadavers after all hair had been clipped and the skin washed. The skin samples are then bathed in 10% glycerin and stored frozen. The glycerin prevents the formation of ice crystals which could possibly damage the keratinized cells and/or the intercellular lipid matrix. After a rapid thawing, the skin is conditioned for 24 hours in Hank's Balanced Salt Solution with 1% antibacterial-antimycotic solution. Then the skin is washed with distilled water. A single skin donor is used for each experiment, and individual sections for use are selected based on integrity of the stratum corneum (visual determination). Selected areas are cut to  $1\text{cm}^2$  using a scalpel.

Tests are conducted using glass diffusion cells placed in temperature-regulated stirring modules. Skin sections are mounted in the cells, and the receptor phase is added. The receptor phase is 50% Hank's Balance Salt Solution (including 1% antibiotic-antimycotic solution) and 50% ethanol. Each diffusion cell has an exposed area of  $0.79\text{cm}^2$  and a receptor capacity of 5ml. Sufficient formulation is applied (750ul) to the surface of the skin to ensure infinite dose conditions, and the diffusion cell is covered with plastic wrap or parafilm to prevent product evaporation. At each sampling time the receptor phase is removed for analysis of drug content. The receptor phase is removed for analysis of drug content. The receptor phase is replenished at each sampling time in order to maintain sink conditions. Preferably 3 to 6 replicates are run with sampling intervals occurring at 1, 2, 4 & 6 hours.

Penetration rate (Flux) is determined as the quantity of drug penetrating a measured area of skin per hour during the 5 hour interval between 1 hour and 6 hours. Generally steady state is reached before 1 hour. Penetration rate is usually expressed as  $\mu\text{g drug per cm}^2\text{ skin per hour}$ .

Ingredients are identified by chemical or CTFA name.

## EXAMPLES

### Example I

An anti-acne composition is made by combining the following components using conventional mixing technology.

Ingredient	(%W/W)
Water, Purified	qs to 100
PPG-14 Butyl Ether	8.0
Polyquaternium-32 and Mineral Oil <sup>1</sup>	2.5
Polyoxyethylene 10 Cetyl Ether	0.75
Salicylic Acid	2.0
Stearyl Alcohol	0.75
Cetyl Alcohol	0.75
85% 5cs Dimethyl Fluid/15% Dimethiconol <sup>2</sup>	0.5
Aluminum Starch Octenyl Succinate	0.5

<sup>1</sup>SalCare<sup>(RTM)</sup> SC92 available from Allied Colloids (Suffolk, VA).

<sup>2</sup>Dow Q2-1403 Fluid available from Dow Corning.

The PPG-14 Butyl ether, salicylic acid, Polyoxyethylene 10 Cetyl Ether, and waxes are combined and heated until the waxes melt (e.g., up to about 75 C) and is mixed using a Lightnin' Mixer with a 3 blade paddle prop at a low speed (100 rpm). Water is added to this mixture. While mixing at a moderate speed (300 rpm), the polyquaternium-32 and mineral oil is added to the mixture. The resulting cream is mixed at moderate speed until uniform.

The composition displays skin penetration of the salicylic acid active with low dermal irritation as well as improved skin feel and residue characteristics together with excellent moisturizing, emolliency, rub-in and absorption characteristics.

Alternatively, the above composition is prepared by substituting the polyquaternium-32 and mineral oil with crosslinked methyl quaternized dimethylaminoethyl methacrylate and mineral oil [i.e. polyquaternium 37 (and) mineral oil (and) PPG-1 trideceth-6], available as SalCare<sup>(RTM)</sup> SC95 from Allied Colloids (Suffolk, VA).

#### Example II

An anti-acne and/or analgesic composition is made by combining the following ingredients utilizing conventional mixing techniques as described above in Example I.

Ingredient	(%W/W)
Water, Purified	qs to 100
Ibuprofen	2.0
Polyquaternium-32 and Mineral Oil <sup>1</sup>	4.0
Polyoxyethylene 10 Cetyl Ether	0.75
Stearyl Alcohol	0.75
Cetyl Alcohol	0.75
85% 5cs Dimethyl Fluid/15% Dimethiconol <sup>2</sup>	0.5
Aluminum Starch Octenyl Succinate	0.5

<sup>1</sup>SalCare<sup>(RTM)</sup> SC92 available from Allied Colloids (Suffolk, VA).

<sup>2</sup>Dow Q2-1403 Fluid available from Dow Corning.

The compositions display skin penetration of the Ibuprofen active as well as improved skin feel and residue characteristics together with excellent moisturizing, emolliency, rub-in and absorption characteristics.

Alternatively, the above composition is prepared by substituting the polyquaternium-32 and mineral oil with crosslinked methyl quaternized dimethylaminoethyl methacrylate (and) mineral oil [i.e. polyquaternium 37 (and) mineral oil (and) PPG-1 trideceth-6], available as SalCare<sup>(RTM)</sup> SC95 from Allied Colloids (Suffolk, VA).

### 5 Example III

An anti-acne cream is made by combining the following components using conventional mixing technology.

Ingredient	(%W/W)
Water, Purified	qs to 100
PPG-14 Butyl Ether	8.0
Crosslinked Methyl Quaternized Dimethylaminoethyl Methacrylate and Mineral Oil <sup>1</sup>	1.5
Steareth-21	0.50
Salicylic Acid	2.0
Stearyl Alcohol	1.5
Cetyl Alcohol	1.5
85% 5cs Dimethyl Fluid/15% Dimethiconol <sup>2</sup>	0.5
Dimethicone 200	0.6
Glycerin	3.0
Citric Acid	0.7
Sodium Citrate	0.3

<sup>1</sup>SalCare<sup>(RTM)</sup> SC95 available from Allied Colloids (Suffolk, VA).

<sup>2</sup>Dow Q2-1403 Fluid available from Dow Corning

In a vessel the cetyl alcohol, stearyl alcohol, steareth-21, and dimethicone 200 are heated to 70°C. In another vessel the PPG-14 butyl ether and salicylic acid are combined and heated to 70°C to dissolve the salicylic acid. This salicylic acid solution is then added with stirring to the cetyl alcohol mixture. Next the crosslinked methyl quaternized dimethylaminoethyl methacrylate polymer is added with stirring to the cetyl alcohol mixture. In a separate vessel, the water, glycerin, citric acid, and sodium citrate are combined and heated with stirring to 70°C. The cetyl alcohol mixture is then added with mixing to the water mixture to form the emulsion. Finally, the dimethyl fluid/dimethicone mixture is added with mixing and the mixture is then cooled to room temperature with mixing.

The resulting cream composition displays skin penetration of the salicylic acid active with low dermal irritation as well as improved skin feel and residue characteristics together with excellent moisturizing, emolliency, rub-in and absorption characteristics.

Alternatively, the above composition is prepared by substituting the methyl quaternized dimethylaminoethyl methacrylate and mineral oil with polyquaternium-32 and mineral oil [i.e. polyquaternium 37 (and) mineral oil (and) PPG-1 trideceth-6], available as SalCare<sup>(RTM)</sup> SC92 from Allied Colloids (Suffolk, VA).

### Example IV

A keratolytic composition for dermatological disorders is made by combining the following ingredients utilizing conventional mixing techniques as described above in Example I.

Ingredient	(%W/W)
Water	qs to 100
Urea	10.0
Polyquaternium-32 and Mineral Oil <sup>1</sup>	4.0
Polyoxyethylene 10 Cetyl Ether	0.75
Stearyl Alcohol	0.75
Cetyl Alcohol	0.75
85% 5cs Dimethyl Fluid/15% Dimethiconol <sup>2</sup>	0.5
Aluminum Starch Octenyl Succinate	0.5

<sup>1</sup>SalCare<sup>(RTM)</sup> SC92 available from Allied Colloids (Suffolk, VA).

<sup>2</sup>Dow Q2-1403 Fluid available from Dow Corning.

The compositions display skin penetration of the Urea active as well as improved skin feel and residue characteristics together with excellent moisturizing, emolliency, rub-in and absorption characteristics.

Alternatively, the above composition is prepared by substituting the polyquaternium-32 and mineral oil with crosslinked methyl quaternized dimethylaminoethyl methacrylate and mineral oil [i.e. polyquaternium 37 (and) mineral oil (and) PPG-1 trideceth-6], available as SalCare<sup>(RTM)</sup> SC95 from Allied Colloids (Suffolk, VA).

#### Example V

A composition for sunless tanning is made by combining the following ingredients utilizing conventional mixing techniques as described above in Example I.

Ingredient	(%W/W)
Water	qs to 100
Polyquaternium-32 and Mineral Oil <sup>1</sup>	3.0
Dihydroxyacetone	3.0
Glycerin	2.0
Polyoxyethylene 10 Cetyl Ether	0.75
Stearyl Alcohol	0.75
Cetyl Alcohol	0.75
85% 5cs Dimethyl Fluid/15% Dimethiconol <sup>2</sup>	0.5
Aluminum Starch Octenyl Succinate	0.5

<sup>1</sup>SalCare<sup>(RTM)</sup> SC92 available from Allied Colloids (Suffolk, VA).

<sup>2</sup>Dow Q2-1403 Fluid available from Dow Corning.

The compositions display improved skin penetration of the dihydroxyacetone as well as improved skin feel and residue characteristics together with excellent moisturizing, emolliency, rub-in and absorption characteristics.

Alternatively, the above composition is prepared by substituting the polyquaternium-32 and mineral oil with crosslinked methyl quaternized dimethylaminoethyl methacrylate and mineral oil [i.e. polyquaternium 37 (and) mineral oil (and) PPG-1 trideceth-6], available as SalCare<sup>(RTM)</sup> SC95 from Allied Colloids (Suffolk, VA).

Example VI

A composition for sunless tanning which also contains sunscreens is made by combining the following ingredients utilizing conventional mixing techniques as described above in Example I.

Ingredient	(%W/W)
Water	qs to 100
Polyquaternium-32 and Mineral Oil <sup>1</sup>	3.0
Dihydroxyacetone	3.0
Octyl Methoxycinnamate	7.5
Octyl Salicylate	1.0
Glycerin	2.0
Polyoxyethylene 10 Cetyl Ether	0.75
Stearyl Alcohol	0.75
Cetyl Alcohol	0.75
85% 5cs Dimethyl Fluid/15% Dimethiconol <sup>2</sup>	0.5
Aluminum Starch Octenyl Succinate	0.5

<sup>1</sup>SalCare<sup>(RTM)</sup> SC92 available from Allied Colloids (Suffolk, VA).

<sup>2</sup>Dow Q2-1403 Fluid available from Dow Corning.

The compositions display improved skin penetration of the dihydroxyacetone as well as improved skin feel and residue characteristics together with excellent moisturizing, emolliency, rub-in and absorption characteristics. This composition also provides protection against the harmful effects of UV radiation.

Alternatively, the above composition is prepared by substituting the polyquaternium-32 and mineral oil with crosslinked methyl quaternized dimethylaminoethyl methacrylate and mineral oil [i.e. polyquaternium 37 (and) mineral oil (and) PPG-1 trideceth-6], available as SalCare<sup>(RTM)</sup> SC95 from Allied Colloids (Suffolk, VA).

Example VII

An anti-acne lotion is made by combining the following components using conventional mixing technology.

Ingredient	(%W/W)
<b>Water Phase</b>	
Water, Purified	qs to 100
Glycerin	3.0
Triethanolamine	0.15
Tetrasodium EDTA	0.02
<b>Oil Phase</b>	
PPG-14 Butyl Ether	8.0
Salicylic Acid	2.0
Crosslinked Methyl Quaternized Dimethylaminoethyl Methacrylate and mineral oil <sup>1</sup>	1.5
Cetyl Alcohol	0.75
Stearyl Alcohol	0.75
Dimethicone <sup>2</sup>	0.60
Steareth-21	0.45
Steareth-2	0.05
<b>Fragrance Phase</b>	
Fragrance	0.08
Cyclomethicone (and) dimethiconol <sup>3</sup>	0.50

<sup>1</sup>SalCare<sup>(RTM)</sup> SC95 [i.e. polyquaternium 37 (and) mineral oil (and) PPG-1 trideceth-6] available from Allied Colloids (Suffolk, VA).

<sup>2</sup>Dow Corning 200/350 centistokes available from Dow Corning.

<sup>3</sup>Dow Q2-1401 Fluid available from Dow Corning.

In an appropriate vessel the water phase ingredients are combined and heated with stirring to about 75°C. In a separate vessel the oil phase ingredients are combined and heated with stirring to about 75°C. The oil phase is then added with stirring to the water phase to form an emulsion. Stirring is continued as the mixture is cooled to about 40°C. The fragrance phase is added and the mixture is cooled to room temperature with stirring.

The composition displays skin penetration of the salicylic acid active with low dermal irritation as well as improved skin feel and residue characteristics together with excellent moisturizing, emolliency, rub-in and absorption characteristics.

In an alternate method of preparation, the crosslinked methyl quaternized dimethylaminoethyl methacrylate and mineral oil is not added to the oil phase, but is instead added after the emulsion has been both formed and subsequently cooled to about 40°C.

Alternatively, the above composition is prepared by substituting the crosslinked methyl quaternized dimethylaminoethyl methacrylate and mineral oil with polyquaternium-32 and mineral oil, available as SalCare<sup>(RTM)</sup> SC92 from Allied Colloids (Suffolk, VA).

#### Example VIII

An anti-acne cream is made by combining the following components using conventional mixing technology.



Ingredient	(%W/W)
<b>Water Phase</b>	
Water, Purified	qs to 100
Glycerin	3.0
Triethanolamine	0.15
Tetrasodium EDTA	0.02
<b>Oil Phase</b>	
PPG-14 Butyl Ether	8.0
Salicylic Acid	2.0
Crosslinked Methyl Quaternized Dimethylaminoethyl Methacrylate and mineral oil <sup>1</sup>	1.5
Cetyl Alcohol	1.50
Stearyl Alcohol	1.50
Dimethicone <sup>2</sup>	0.60
Steareth-21	0.45
Steareth-2	0.05
<b>Fragrance Phase</b>	
Fragrance	0.08
Cyclomethicone (and) dimethiconol <sup>3</sup>	0.50

<sup>1</sup>SalCare<sup>(RTM)</sup> SC95 [i.e. polyquaternium 37 (and) mineral oil (and) PPG-1 trideceth-6] available from Allied Colloids (Suffolk, VA).

<sup>2</sup>Dow Corning 200/350 centistokes available from Dow Corning.

<sup>3</sup>Dow Q2-1401 Fluid available from Dow Corning.

In an appropriate vessel the water phase ingredients are combined and heated with stirring to about 75°C. In a separate vessel the oil phase ingredients are combined and heated with stirring to about 75°C. The oil phase is then added with stirring to the water phase to form an emulsion. Stirring is continued as the mixture is cooled to about 40°C. The fragrance phase is added and the mixture is cooled to room temperature with stirring.

The composition displays skin penetration of the salicylic acid active with low dermal irritation as well as improved skin feel and residue characteristics together with excellent moisturizing, emolliency, rub-in and absorption characteristics.

In an alternate method of preparation, the crosslinked methyl quaternized dimethylaminoethyl methacrylate and mineral oil is not added to the oil phase, but is instead added after the emulsion has been both formed and subsequently cooled to about 40°C.

Alternatively, the above composition is prepared by substituting the crosslinked methyl quaternized dimethylaminoethyl methacrylate and mineral oil with polyquaternium-32 and mineral oil, available as SalCare<sup>(RTM)</sup> SC92 from Allied Colloids (Suffolk, VA).

#### EXAMPLE IX

A salicylic acid containing cream is made by combining the following components using conventional mixing technology.

Ingredient	(%W/W)
<b>Water Phase</b>	
Water, Purified	qs to 100
Glycerin	3.0
Triethanolamine	0.15
Tetrasodium EDTA	0.02
<b>Oil Phase</b>	
PPG-14 Butyl Ether	16.0
Salicylic Acid	4.0
Crosslinked Methyl Quaternized Dimethylaminoethyl Methacrylate and mineral oil <sup>1</sup>	3.0
Cetyl Alcohol	0.75
Stearyl Alcohol	0.75
Dimethicone <sup>2</sup>	0.60
Steareth-21	0.45
Steareth-2	0.05
<b>Fragrance Phase</b>	
Fragrance	0.08
Cyclomethicone (and) dimethiconol <sup>3</sup>	0.50

<sup>1</sup>SalCare<sup>(RTM)</sup> SC95 [i.e. polyquaternium 37 (and) mineral oil (and) PPG-1 trideceth-6] available from Allied Colloids (Suffolk, VA).

<sup>2</sup>Dow Corning 200/350 centistokes available from Dow Corning.

<sup>3</sup>Dow Q2-1401 Fluid available from Dow Corning.

In an appropriate vessel the water phase ingredients are combined and heated with stirring to about 75°C. In a separate vessel the oil phase ingredients are combined and heated with stirring to about 75°C. The oil phase is then added with stirring to the water phase to form an emulsion. Stirring is continued as the mixture is cooled to about 40°C. The fragrance phase is added and the mixture is cooled to room temperature with stirring.

The composition displays skin penetration of the salicylic acid active with low dermal irritation as well as improved skin feel and residue characteristics together with excellent moisturizing, emolliency, rub-in and absorption characteristics.

In an alternate method of preparation, the crosslinked methyl quaternized-dimethylaminoethyl methacrylate and mineral oil is not added to the oil phase, but is instead added after the emulsion has been both formed and subsequently cooled to about 40°C.

Alternatively, the above composition is prepared by substituting the crosslinked methyl quaternized dimethylaminoethyl methacrylate and mineral oil with polyquaternium-32 and mineral oil, available as SalCare<sup>(RTM)</sup> SC92 from Allied Colloids (Suffolk, VA).


## Claims

1. A topical pharmaceutical composition having enhanced penetration through the skin comprising:

- (a) a safe and effective amount of a pharmaceutical active;
- (b) from 0.1% to 10.0% of a high molecular weight crosslinked cationic polymer of the formula: (A)<sub>l</sub>(B)<sub>m</sub>(C)<sub>n</sub> wherein (A) is a dialkylaminoalkyl acrylate monomer or its quaternary ammonium or acid addition salt, (B) is a dialkylaminoalkyl methacrylate monomer or its quaternary ammonium or acid addition salt, (C) is acrylamide, l is an integer of 0 or greater, m is an integer of 1 or greater, and n is an integer of 0 or greater, wherein said polymer contains a crosslinking agent;

(c) from 0.05% to 5% of a high HLB non-ionic surfactant selected from ethoxylated alcohols having from 8 to 22 carbons and from 6 to 30 ethylene oxide groups; and

(d) from 0.1% to 25% of an alkoxyated ether of the formula:



wherein R is straight or branched chain C<sub>1</sub> to C<sub>18</sub>, and X is methyl or ethyl, and wherein n has an average value of from 5 to 20;

wherein the composition is substantially free of ethanol.


2. The composition of Claim 1 wherein the crosslinking agent is selected from methylene bisacrylamide, ethylene glycol di-(meth)acrylate, di-(meth)acrylamide, cyanomethylacrylate, vinyloxyethylacrylate, vinyloxyethylmethacrylate, allyl pentaerythritol, trimethylolpropane diallylether, allyl sucrose, butadiene, isoprene, divinyl benzene, divinyl naphthalene, allyl acrylate, and mixtures thereof; preferably methylene bisacrylamide.

3. The composition of Claim 2 wherein said high HLB nonionic surfactant is selected from ceteth-10, steareth-21, and mixtures thereof, and wherein said alkoxyated ether is selected from PPG-14 butyl ether, PPG-15 stearyl ether, and mixtures thereof.

4. The composition of Claim 3 wherein said pharmaceutical active is selected from anti-acne drugs, anti-wrinkle actives, anti-skin atrophy actives, non-steroidal anti-inflammatory drugs, steroidal anti-inflammatory drugs, sunless tanning agents, sunscreen agents, wound healing agents, skin bleaching or lightening agents, antihistaminic drugs, antitussive drugs, antipruritic drugs, anticholinergic drugs, anti-emetic and antinauseant drugs, anorexic drugs, central stimulant drugs, antiarrhythmic drugs, B-adrenergic blocker drugs, cardiotonic drugs, antihypertensive drugs, diuretic drugs, vasodilator drugs, vasoconstrictor drugs, anti-ulcer drugs, anesthetic drugs, antidepressant drugs, tranquilizer and sedative drugs, antipsychotic drugs, antimicrobial drugs, antineoplastic drugs, antimalarial drugs, muscle relaxant drugs, anti spasmodic drugs, antidiarrheal drugs and bone-active drugs and mixtures thereof; preferably wherein said antihistaminic drug is selected from chlorpheniramine maleate, chlorpheniramine tannate, triprolidine hydrochloride, triprolidine oxalate, diphenhydramine hydrochloride, diphenhydramine ascorbate, diphenhydramine citrate, doxylamine succinate, pyrilamine maleate, pyrilamine hydrochloride, pyrilamine tannate, phenindamine tartrate, promethazine hydrochloride, cyproheptadine hydrochloride, azatadine maleate, clemastine fumarate, carbinoxamine maleate, carbinoxamine hydrochloride, tripeleonnamine hydrochloride, tripeleonnamine citrate, dexchlorpheniramine maleate, brompheniramine maleate and chlorcyclizine hydrochloride and mixtures thereof; wherein said antitussive drug is selected from the group consisting of dextromethorphan hydrobromide, carbetapentane citrate, codeine phosphate and codeine N-oxide hydrochloride and mixtures thereof; wherein said anticholinergic drug is selected from the group consisting of scopolamine hydrobromide, scopolamine hydrochloride, atropine sulfate, atropine mucate, homatropine hydrobromide and homatropine hydrochloride and mixtures thereof; wherein said anti-emetic or antinauseant drug is selected from cyclizine hydrochloride, meclizine hydrochloride, chlorpromazine hydrochloride and chlorpromazine maleate and mixtures thereof; wherein said anorexic drug is selected from benzphetamine hydrochloride, phentermine hydrochloride, chlorphentermine hydrochloride and fenfluramine hydrochloride and mixtures thereof; wherein said antimicrobial drug is selected from  $\beta$ -lactam drugs, quinolone drugs, ciprofloxacin, norfloxacin, tetracycline, erythromycin, hexamidine isethionate, amikacin, triclosan, doxycycline, capreomycin, chlorhexidine, chlortetracycline, oxytetracycline, clindamycin, ethambutol, metronidazole, pentamidine, gentamicin, kanamycin, lineomycin, methacycline, methenamine, minocycline, neomycin, netilmicin, paromomycin, streptomycin, tobramycin, miconazole and amantadine, pharmaceutically-acceptable salts thereof and mixtures thereof; wherein said antiarrhythmic drug is selected from propranolol hydrochloride, procainamide hydrochloride, quinidine sulfate and quinidine gluconate and mixtures thereof; wherein said antihypertensive drug is selected from enalapril maleate, clonidine hydrochloride, hydralazine hydrochloride and hydralazine sulfate and mixtures thereof; wherein said anesthetic or antipruritic drug is selected from lidocaine hydrochloride, bupivacaine hydrochloride, chlorprocaine hydrochloride, dibucaine hydrochloride, etidocaine hydrochloride, mepivacaine hydrochloride, tetracaine hydrochloride, dyclonine hydrochloride and hexylcaine hydrochloride and mixtures thereof; wherein said bone-active drug is selected from 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, 3-amino-1-hydroxypropane-1,1-diphosphonic acid, octahydro-1-pyridine-6,6-diphosphonic acid, 2-(2'-piperidinyl)ethane-1,1-diphosphonic acid; 2-(3'-piperidinyl)-ethane-1,1-diphosphonic acid; 2-(2'-piperidinyl)-1-hydroxyethane-1,1-diphosphonic acid; 2-(3'-piperidinyl)-1-hydroxyethane-1,1-diphosphonic acid; N-(2'-(3'-

methyl)-piperidinylidene)-amino-methane diphosphonic acid; N-(2'-(1',3'-diazinyliidene))-aminomethane diphosphonic acid; and N-(2-(3-methyl-piperidinyliidene))-aminomethanephosphonomethylphosphinic acid, or esters thereof and mixtures thereof; wherein said non-steroidal anti-inflammatory drug is selected from propionic acid derivatives, acetic acid derivatives, fenamic acid derivatives, biphenylcarboxylic acid derivatives, and oxicams and mixtures thereof; and wherein said non-steroidal anti-inflammatory drug is a propionic acid derivatives selected from aspirin, acetaminophen, ibuprofen, naproxen, benoxaprofen, flurbiprofen, fenoprofen, fenbufen, ketoprofen, indoprofen, pirofen, carprofen, oxaprozin, pranoprofen, miroprofen, tioxaprofen, suprofen, alminoprofen, tiaprofenic acid, fluprofen and buclocic acid and mixtures thereof.

5. The composition of Claim 3 wherein said pharmaceutical active is an anti-acne drug selected from salicylic acid, sulfur, resorcinol, N-acetylcysteine, octopirox, retinoic acid and its derivatives, benzoyl peroxide, erythromycin, zinc, tetracyclin, azelaic acid and its derivatives, phenoxy ethanol and phenoxy propanol, ethylacetate, clindamycin and medocycline, flavinoids, lactic acid, glycolic acid, pyruvic acid, urea, scymnol sulfate and its derivatives, deoxycholate and cholate and mixtures thereof, or an anti-wrinkle active or anti-skin atrophy active selected from glycolic acid, lactic acid, salicylic acid, and mixtures thereof.
6. The composition of Claim 3 wherein said pharmaceutical active is selected from salicylic acid, glycolic acid, lactic acid, and mixtures thereof.
7. The composition of Claim 5 wherein the amount of (C) in the cationic polymer is from 50% to 90% molar; preferably wherein l in the cationic polymer is zero and the ratio of (B):(C) is from 45:55 to 55:45; more preferably wherein both l and n are zero in the cationic polymer.
8. The composition of Claim 7 which further comprises from 3% to 5% glycerin, and from 1% to 10% of an emollient selected from volatile silicone oils, non-volatile silicone oils, highly branched hydrocarbons, and mixtures thereof.
9. The composition of Claim 3 wherein said drug active is a sunless tanning agent selected from dihydroxyacetone, indole derivatives and mixtures thereof; preferably wherein said composition further comprises a sunscreen active selected from 2-ethylhexyl p-methoxycinnamate, 2-ethylhexyl N,N-dimethyl-p-aminobenzoate, p-aminobenzoic acid, 2-phenylbenzimidazole-5-sulfonic acid, octocrylene, oxybenzone, homomenthyl salicylate, octyl salicylate, 4,4'-methoxy-t-butylidibenzoylmethane, 4-isopropyl dibenzoylmethane, 3-benzylidene camphor, 3-(4-methylbenzylidene) camphor, titanium dioxide, zinc oxide, silica, iron oxide, and mixtures thereof.
10. A topical pharmaceutical composition having enhanced penetration through the skin comprising:
  - (a) a safe and effective amount of a pharmaceutical active;
  - (b) from 0.1% to 10.0% of a high molecular weight crosslinked cationic polymer of the formula:  $(A)_l(B)_m(C)_n$  wherein (A) is a dialkylaminoalkyl acrylate monomer or its quaternary ammonium or acid addition salt, (B) is a dialkylaminoalkyl methacrylate monomer or its quaternary ammonium or acid addition salt, (C) is acrylamide, l is an integer of 0 or greater, m is an integer of 1 or greater, and n is an integer of 0 or greater, wherein said polymer contains a crosslinking agent;
  - (c) from 0.05% to 5% of a high HLB non-ionic surfactant selected from ethoxylated alcohols having from 8 to 22 carbons and from 6 to 30 ethylene oxide groups; and
  - (d) from 0.01% to 5% of a low HLB non-ionic surfactant; and
  - (e) from 0.1% to 25% of an alkoxyated ether of the formula:



wherein R is straight or branched chain  $\text{C}_1$  to  $\text{C}_{18}$ , and X is methyl or ethyl, and wherein n has an average value of from 5 to 20;

wherein the composition is substantially free of ethanol.

11. The composition of Claim 10 wherein the crosslinking agent is selected from methylene bisacrylamide, ethylene glycol di-(meth)acrylate; di-(meth)acrylamide, cyanomethylacrylate, vinyloxyethylacrylate, vinyloxyethylmethacrylate, allyl pentaerythritol, trimethylolpropane diallyl ether, allyl sucrose, butadiene, isoprene, divinyl benzene, divi-


nyl naphthalene, allyl acrylate, and mixtures thereof; preferably wherein said crosslinking agent is methylene bisacrylamide.

12. The composition of Claim 11 wherein said high HLB nonionic surfactant is selected from ceteth-10, steareth-21, and mixtures thereof, and wherein said alkoxylated ether is selected from PPG-14 butyl ether, PPG-15 stearyl ether, and mixtures thereof.
13. The composition of Claim 12 wherein said low HLB nonionic surfactant is selected from ethoxylated alcohols having from 8 to 22 carbons and from 1 to 5 ethylene oxide groups; preferably wherein said low HLB nonionic surfactant is selected from steareth-2, ceteth-2, laureth-2, oleth-2, and mixtures thereof; and more preferably wherein said low HLB nonionic surfactant is steareth-2.
14. The composition of Claim 13 wherein said pharmaceutical active is selected from anti-acne drugs, anti-wrinkle actives, anti-skin atrophy actives, non-steroidal anti-inflammatory drugs, steroidal anti-inflammatory drugs, sunless tanning agents, sunscreen agents, wound healing agents, skin bleaching or lightening agents, antihistaminic drugs, antitussive drugs, antipruritic drugs, anticholinergic drugs, anti-emetic and antinauseant drugs, anorexic drugs, central stimulant drugs, antiarrhythmic drugs, B-adrenergic blocker drugs, cardiogenic drugs, antihypertensive drugs, diuretic drugs, vasodilator drugs, vasoconstrictor drugs, anti-ulcer drugs, anesthetic drugs, antidepressant drugs, tranquilizer and sedative drugs, antipsychotic drugs, antimicrobial drugs, antineoplastic drugs, antimalarial drugs, muscle relaxant drugs, antispasmodic drugs, antidiarrheal drugs and bone-active drugs and mixtures thereof.
15. The composition of Claim 13 wherein said pharmaceutical active is an anti-acne drug selected from salicylic acid, sulfur, resorcinol, N-acetylcysteine, octopirox, retinoic acid and its derivatives, benzoyl peroxide, erythromycin, zinc, tetracyclin, azelaic acid and its derivatives, phenoxy ethanol and phenoxy propanol, ethylacetate, clindamycin and medocycline, flavinoids, lactic acid, glycolic acid, pyruvic acid, urea, scymnol sulfate and its derivatives, deoxycholate and cholate and mixtures thereof, or an anti-wrinkle active or anti-skin atrophy active selected from glycolic acid, lactic acid, salicylic acid, and mixtures thereof.
16. The composition of Claim 13 wherein said pharmaceutical active is selected from salicylic acid, lactic acid, glycolic acid, and mixtures thereof.
17. The composition of Claim 13 wherein the amount of (C) in the cationic polymer is from 50% to 90% molar; preferably wherein I in the cationic polymer is zero and the ratio of (B):(C) is from 45:55 to 55:45; and more preferably wherein both I and n are zero in the cationic polymer.
18. The composition of Claim 17 which further comprises from 3% to 5% glycerin and from 1% to 10% of an emollient selected from the group consisting of volatile silicone oils, non-volatile silicone oils, highly branched hydrocarbons, and mixtures thereof.
19. The composition of Claim 13 wherein said drug active is a sunless tanning agent selected from dihydroxyacetone, indole derivatives and mixtures thereof; preferably wherein said composition further comprises a sunscreen active selected from 2-ethylhexyl p-methoxycinnamate, 2-ethylhexyl N,N-dimethyl-p-aminobenzoate, p-aminobenzoic acid, 2-phenylbenzimidazole-5-sulfonic acid, octocrylene, oxybenzone, homomenthyl salicylate, octyl salicylate, 4,4'-methoxy-1-butylidibenzoylmethane, 4-isopropyl dibenzoylmethane, 3-benzylidene camphor, 3-(4-methylbenzylidene) camphor, titanium dioxide, zinc oxide, silica, iron oxide, and mixtures thereof.

#### Patentansprüche

1. Topische pharmazeutische Zusammensetzung mit gesteigerter Penetration durch die Haut, umfassend:
  - (a) eine sichere und wirksame Menge eines aktiven Pharmazeutikums;
  - (b) von 0,1 % bis 10,0 % eines hochmolekularen vernetzten kationischen Polymers mit der Formel:  $(A)_l(B)_m(C)_n$ , worin (A) ein Dialkylaminoalkylacrylatmonomer oder dessen quaternäres Ammonium-oder Säureadditionssalz ist, (B) ein Dialkylaminoalkylmethacrylatmonomer oder dessen quaternäres Ammonium-oder Säureadditionssalz ist, (C) Acrylamid bedeutet, l eine ganze Zahl von 0 oder darüber ist, m eine ganze Zahl von 1 oder darüber ist und n eine ganze Zahl von 0 oder darüber ist, wobei dieses Polymer ein Vernetzungsmittel enthält;
  - (c) von 0,05 % bis 5 % eines nichtionischen grenzflächenaktiven Mittels mit hohem HLB-Wert, ausgewählt

unter ethoxylierten Alkoholen mit 8 bis 22 Kohlenstoffatomen und 6 bis 30 Ethylenoxidgruppen; und  
(d) von 0,1 % bis 25 % eines alkoxylierten Ethers mit der Formel:



5

10 worin R gerades oder verzweigt-kettiges C<sub>1</sub> bis C<sub>18</sub> bedeutet und X für Methyl oder Ethyl steht und n einen Mittelwert von 5 bis 20 aufweist;

worin die Zusammensetzung im wesentlichen frei von Ethanol ist.

- 15 2. Zusammensetzung nach Anspruch 1, worin das Vernetzungsmittel ausgewählt ist unter Methylenbisacrylamid, Ethylenglycoldi-(meth)acrylat, Di-(meth)acrylamid, Cyanomethylacrylat, Vinyloxyethylacrylat, Vinyloxyethylmethacrylat, Allylpentaerythrit, Trimethylolpropandiallyther, Allylsaccharose, Butadien, Isopren, Divinylbenzol, Divinylnaphtalin, Allylacrylat und Gemischen hievon; vorzugsweise Methylenbisacrylamid.
- 20 3. Zusammensetzung nach Anspruch 2, worin das nichtionische grenzflächenaktive Mittel mit hohem HLB-Wert ausgewählt ist unter Ceteth-10, Steareth-21 und Gemischen hievon und worin der alkoxylierte Ether ausgewählt ist unter PPG-14-butylether, PPG-15-stearylether und Gemischen hievon.
- 25 4. Zusammensetzung nach Anspruch 3, worin das aktive Pharmazeutikum ausgewählt ist unter Anti-Aknemitteln, Anti-Faltenmitteln, Anti-Hautatrophiemitteln, nichtsteroiden entzündungsbemmenden Mitteln, steroiden entzündungshemmenden Mitteln, sonnenfreien Bräunungsmitteln, Sonnenschutzmitteln, Wundheilmitteln, Hautbleich- oder -aufhellungsmitteln, Antihistaminika, Antitussiva, Antipruritusmitteln, Anticholinergika, Antiemetika und Antinauseantmitteln, Anorektika, Zentralstimulantien, Antiarrhythmika,  $\beta$ -adrenergen Blockern, Cardiotonika, Antihypertensiva, Diuretika, Vasodilatoren, Vasoconstrictoren, Antiulzera, Anästhetika, Antidepressiva, Tranquilizern und Sedativa, Antipsychotika, antimikrobiellen Mitteln, antineoplastischen Mitteln, Antimalariamitteln, Muskelrelaxantien, Antispasmodika, Antidiarrhoemitteln und knochenaktiven Mitteln und Gemischen hievon; wobei vorzugsweise das Antihistaminikum ausgewählt ist unter Chlorpheniraminmaleat, Chlorpheniraminantannat, Tripolidinhydrochlorid, Tripolidinoxalat, Diphenhydraminhydrochlorid, Diphenhydraminascorbat, Diphenhydramincitrat, Doxylamin-30 succinat, Pyrilaminmaleat, Pyrilaminhydrochlorid, Pyrilaminantannat, Phenindamintartrat, Promethazinhydrochlorid, Cyproheptadinhydrochlorid, Azatadinmaleat, Clemastinmaleat, Carbinoxaminmaleat, Carbinoxaminhydrochlorid, Tripelennaminhydrochlorid, Tripelennaminantannat, Dexchlorpheniraminmaleat, Brompheniraminmaleat und Chlorcyclizinhydrochlorid und Gemischen hievon; worin das Antitussivum aus der aus Dextromethorphanhydrobromid, Carbetapentancitrat, Codeinphosphat und Codein-N-oxid-hydrochlorid und deren Gemischen bestehenden Gruppe ausgewählt ist; worin das Anticholinergikum aus der aus Scopolaminhydrobromid, Scopolaminhydrochlorid, Atropinsulfat, Atropinmucal, Homatropinhydrobromid und Homatropinhydrochlorid und Gemischen hievon bestehenden Gruppe ausgewählt ist; worin das Antiemetikum oder das Antinauseamittel ausgewählt ist unter Cyclizinhydrochlorid, Meclizinhydrochlorid, Chlorpromazinhydrochlorid und Chlorpromazinmaleat und Gemischen hievon; worin das Anorektikum ausgewählt ist unter Benzphetaminhydrochlorid, Phenterminhydrochlorid, Chlorphenterminhydrochlorid und Fenfluraminhydrochlorid und Gemischen hievon; worin das antimikrobielle Mittel ausgewählt ist unter  $\beta$ -Lactammitteln, -Chinolonen, Ciprofloxacin, Norfloxacin, Tetracyclin, Erythromycin, Hexamidinisethionat, Amikacin, Triclosan, Doxycyclin, Capreomycin, Chlorhexidin, Chlortetracyclin, Oxytetracyclin, Clindamycin, Ethambutol, Metronidazol, Pentamidin, Gentamicin, Kanamycin, Lineomycin, Methacyclin, Methenamin, Minocyclin, Neomycin, Netilmicin, Paromomycin, Streptomycin, Tobramycin, Miconazol und Amanfadin, pharmazeutisch annehmbaren Salzen hievon und Gemischen hievon; worin das Antiarrhythmikum ausgewählt ist unter Propranololhydrochlorid, Procainamidhydrochlorid, Chinidinsulfat und Chinidingluconat und Gemischen hievon; worin das Antihypertensivum ausgewählt ist unter Enalaprilmaleat, Clonidinhydrochlorid, Hydralazinhydrochlorid und Hydralazinsulfat und Gemischen hievon; worin das Anästhetikum oder Antipruritusmittel ausgewählt ist unter Lidocainhydrochlorid, Bupivacainhydrochlorid, Chlorprocainhydrochlorid, Dibucainhydrochlorid, Etidocainhydrochlorid, Mepivacainhydrochlorid, Tetracainhydrochlorid, Dycloninhydrochlorid und Hexylcainhydrochlorid und Gemischen hievon; worin das knochenaktive Mittel ausgewählt ist unter 6-Amino-1-hydroxyhexan-1,1-diphosphonsäure, 3-Amino-1-hydroxypropan-1,1-diphosphonsäure, Octahydro-1-pyridin-6,6-diphosphonsäure; 2-(2'-Piperidinyl)ethan-1,1-diphosphonsäure; 2-(3'-Piperidinyl)-ethan-1,1-diphosphonsäure; 2-(2'-Piperidinyl)-1-hydroxyethan-1,1-diphosphonsäure; 2-(3'-Piperidinyl)-1-hydroxyethan-1,1-diphosphonsäure; N-(2'-(3'-Methyl)-piperidinyliden)-aminomethandiphosphonsäure; N-(2'-(1',3'-Diazinyliden))-aminomethandiphosphonsäure; und N-(2-(3-Methyl-
- 55

- piperidinylden))-aminomethanphosphonomethylphosphinsäure, oder deren Estern und Gemischen hievon; worin das nicht-steroidale entzündungshemmende Mittel ausgewählt ist unter Propionsäurederivaten, Essigsäurederivaten, Fenaminsäurederivaten, Biphenylcarbonsäurederivaten und Oxicamen und Gemischen hievon; und worin das nichtsteroidale entzündungshemmende Mittel ein Propionsäurederivat ist, ausgewählt unter Aspirin, Acetaminophen, Ibuprofen, Naproxen, Benoxaprofen, Flurbiprofen, Fenoprofen, Fenbuten, Ketoprofen, Indoprofen, Pirprofen, Carprofen, Oxaprozin, Pranoprofen, Miroprofen, Tioxaprofen, Suprofen, Alminoprofen, Tiaprofensäure, Fluprofen und Bucloxinsäure und Gemischen hievon.
- 5 5. Zusammensetzung nach Anspruch 3, worin das aktive Pharmazeutikum ein Anti-Aknemittel ist, ausgewählt unter Salicylsäure, Schwefel, Resorcin, N-Acetylcystein, Octopirox, Retinoesäure und deren Derivaten, Benzoylperoxid, Erythromycin, Zink, Tetracyclin, Azelainsäure und deren Derivaten, Phenoxyethanol und Phenoxypropanol, Ethylacetat, Clindamycin und Meclocylin, Flavonoiden, Milchsäure, Glycolsäure, Brenztraubensäure, Harnstoff, Scymnolsulfat und dessen Derivaten, Deoxycholat und Cholat und Gemischen hievon, oder ein Anti-Faltenwirkstoff oder Anti-Hautatrophie-Wirkstoff ist, ausgewählt unter Glycolsäure, Milchsäure, Salicylsäure und Gemischen hievon.
- 10 6. Zusammensetzung nach Anspruch 3, worin das wirksame Pharmazeutikum ausgewählt ist unter Salicylsäure, Glycolsäure, Milchsäure und Gemischen hievon.
7. Zusammensetzung nach Anspruch 5, worin die Menge (C) im kationischen Polymer 50 bis 90 Mol% beträgt; worin vorzugsweise I im kationischen Polymer den Wert Null hat und das Verhältnis (B):(C) von 45:55 bis 55:45 beträgt; worin stärker bevorzugt sowohl I als auch n im kationischen Polymer den Wert Null aufweisen.
- 15 8. Zusammensetzung nach Anspruch 7, die zusätzlich 3 bis 5 % Glycerin und 1 bis 10 % eines unter flüchtigen Silikonölen, nichtflüchtigen Silikonölen, stark verzweigten Kohlenwasserstoffen und Gemischen hievon ausgewählten Erweichungsmittels umfaßt.
- 25 9. Zusammensetzung nach Anspruch 3, worin der aktive Wirkstoff ein sonnenfreies Bräunungsmittel, ausgewählt unter Dihydroxyaceton, Indolderivaten und Gemischen hievon, ist; worin vorzugsweise diese Zusammensetzung weiterhin ein aktives Sonnenschutzmittel umfaßt, ausgewählt unter 2-Ethylhexyl-p-methoxycinnamat, 2-Ethylhexyl-N,N-dimethyl-p-aminobenzoat, p-Aminobenzoessäure, 2-Phenylbenzimidazol-5-sulfonsäure, Octocrylen, Oxybenzon, Homomenthylsalicylat, Octylsalicylat, 4,4'-Methoxy-t-butyldibenzoylmethan, 4-Isopropyl-dibenzoylmethan, 3-Benzyliden-campher, 3-(4-Methylbenzyliden)campher, Titandioxid, Zinkoxid, Siliciumdioxid, Eisenoxid und Gemischen hievon.
- 30 10. Topische pharmazeutische Zusammensetzung mit gesteigerter Penetration durch die Haut, umfassend:
- (a) eine sichere und wirksame Menge eines aktiven Pharmazeutikums;
- (b) von 0,1 % bis 10,0 % eines hochmolekularen vernetzten kationischen Polymers mit der Formel:  $(A)_l(B)_m(C)_n$ , worin (A) ein Dialkylaminoalkylacrylatmonomer oder dessen quaternäres Ammonium- oder Säureadditionssalz ist, (B) ein Dialkylaminoalkylmethacrylatmonomer oder dessen quaternäres Ammonium- oder Säureadditionssalz ist, (C) Acrylamid bedeutet, l eine ganze Zahl von 0 oder darüber ist, m eine ganze Zahl von 1 oder darüber ist und n eine ganze Zahl von 0 oder darüber ist, wobei dieses Polymer ein Vernetzungsmittel enthält;
- (c) von 0,05 % bis 5 % eines nichtionischen grenzflächenaktiven Mittels mit hohem HLB-Wert, ausgewählt unter ethoxylierten Alkoholen mit 8 bis 22 Kohlenstoffatomen und 6 bis 30 Ethylenoxidgruppen; und
- (d) von 0,01 % bis 5 % eines nichtionischen grenzflächenaktiven Mittels mit niedrigem HLB-Wert; und
- (e) von 0,1 % bis 25 % eines alkoxylierten Ethers mit der Formel:



55 worin R gerades oder verzweigt-kettiges  $\text{C}_1$  bis  $\text{C}_{18}$  bedeutet und X für Methyl oder Ethyl steht und n einen Mittelwert von 5 bis 20 aufweist;

worin die Zusammensetzung im wesentlichen frei von Ethanol ist.



11. Zusammensetzung nach Anspruch 10, worin das Vernetzungsmittel ausgewählt ist unter Methylenbisacrylamid, Ethylenglycoldi-(meth)acrylat, Di-(meth)acrylamid, Cyanomethylacrylat, Vinyloxyethylacrylat, Vinyloxyethylmethacrylat, Allylpentaerythrit, Trimethylolpropandiallyther, Allylsaccharose, Butadien, Isopren, Divinylbenzol, Divinylnaphtalin, Allylacrlat und Gemischen hievon; worin das Vernetzungsmittel vorzugsweise Methylenbisacrylamid ist.
12. Zusammensetzung nach Anspruch 11, worin das nichtionische grenzflächenaktive Mittel mit hohem HLB-Wert ausgewählt ist unter Ceteth-10, Steareth-21 und Gemischen hievon und worin der alkoxylierte Ether ausgewählt ist unter PPG-14-butylether, PPG-15-stearylether und Gemischen hievon.
13. Zusammensetzung nach Anspruch 12, worin das nichtionische grenzflächenaktive Mittel mit niedrigem HLB-Wert unter ethoxlierten Alkoholen mit 8 bis 22 Kohlenstoffomen und 1 bis 5 Ethylenoxidgruppen ausgewählt ist; worin vorzugsweise dieses nichtionische grenzflächenaktive Mittel mit niedrigem HLB-Wert unter Steareth-2, Ceteth-2, Laureth-2, Oleth-2 und Gemischen hievon ausgewählt ist; und stärker bevorzugt das nichtionische grenzflächenaktive Mittel mit niedrigem HLB-Wert Steareth-2 ist.
14. Zusammensetzung nach Anspruch 13, worin das aktive Pharmazeutikum ausgewählt ist unter Anti-Aknemitteln, Anti-Faltenmitteln, Anti-Hautatrophiemitteln, nichtsteroiden entzündungshemmenden Mitteln, steroiden entzündungshemmenden Mitteln, sonnenfreien Bräunungsmitteln, Sonnenschutzmitteln, Wundheilmitteln, Hautbleich- oder -aufhellungsmitteln, Antihistaminika, Antitussiva, Antipruritusmitteln, Anticholinergika, Antiemetika und Antinauseantmitteln, Anorektika, Zentralstimulantien, Antiarrhythmika,  $\beta$ -adrenergen Blockern, Cardiotonika, Antihypertensiva, Diuretika, Vasodilatoren, Vasoconstrictoren, Antiulzera, Anästhetika, Antidepressiva, Tranquilizern und Sedativa, Antipsychotika, antimikrobiellen Mitteln, antineoplastischen Mitteln, Antimalariamitteln, Muskelrelaxantien, Antispasmodika, Antidiarrhoemitteln und knochenaktiven Mitteln und Gemischen hievon.
15. Zusammensetzung nach Anspruch 13, worin das aktive Pharmazeutikum ein Anti-Aknemittel ist, ausgewählt unter Salicylsäure, Schwefel, Resorcin, N-Acetylcystein, Octopirox, Retinoesäure und deren Derivaten, Benzoylperoxid, Erythromycin, Zink, Tetracyclin, Azelainsäure und deren Derivaten, Phenoxyethanol und Phenoxypropanol, Ethylacetat, Clindamycin und Meclocyclin, Flavinoiden, Milchsäure, Glycolsäure, Brenztraubensäure, Harnstoff, Scymnolsulfat und dessen Derivaten, Deoxycholat und Cholat und Gemischen hievon, oder ein Anti-Faltenwirkstoff oder Anti-Hautatrophie-Wirkstoff ist, ausgewählt unter Glycolsäure, Milchsäure, Salicylsäure und Gemischen hievon.
16. Zusammensetzung nach Anspruch 13, worin das wirksame Pharmazeutikum ausgewählt ist unter Salicylsäure, Glycolsäure, Milchsäure und Gemischen hievon.
17. Zusammensetzung nach Anspruch 13, worin die Menge (C) im kationischen Polymer 50 bis 90 Mol% beträgt; worin vorzugsweise I im kationischen Polymer den Wert Null hat und das Verhältnis (B):(C) von 45:55 bis 55:45 beträgt; worin stärker bevorzugt sowohl I als auch n im kationischen Polymer den Wert Null aufweisen.
18. Zusammensetzung nach Anspruch 17, die zusätzlich 3 bis 5 % Glycerin und 1 bis 10 % eines unter flüchtigen Silikonölen, nichtflüchtigen Silikonölen, stark verzweigten Kohlenwasserstoffen und Gemischen hievon ausgewählten Erweichungsmittels umfaßt.
19. Zusammensetzung nach Anspruch 13, worin der aktive Wirkstoff ein sonnenfreies Bräunungsmittel, ausgewählt unter Dihydroxyaceton, Indolderivaten und Gemischen hievon, ist; worin vorzugsweise diese Zusammensetzung weiterhin ein aktives Sonnenschutzmittel umfaßt, ausgewählt unter 2-Ethylhexyl-p-methoxycinnamat, 2-Ethylhexyl-N,N-dimethyl-p-aminobenzoat, p-Aminobenzoessäure, 2-Phenylbenzimidazol-5-sulfonsäure, Octocrylen, Oxybenzon, Homomenthylsalicylat, Octylsalicylat, 4,4'-Methoxy-t-butylidibenzoylmethan, 4-Isopropyl-dibenzoylmethan, 3-Benzyliden-campher, 3-(4-Methylbenzyliden)campher, Titandioxid, Zinkoxid, Siliciumdioxid, Eisenoxid und Gemischen hievon.


## Revendications

1. Composition pharmaceutique à usage local présentant une pénétration de la peau améliorée, comprenant:
- (a) une quantité sans danger et efficace d'un principe pharmaceutique actif;
- (b) de 0,1% à 10,0% d'un polymère cationique réticulé de masse moléculaire élevée, de formule:  $(A)_l(B)_m(C)_n$ , dans laquelle (A) est un monomère acrylate de dialkylaminoalkyle, ou son sel d'ammonium quaternaire ou son sel d'addition d'acide, (B) est un monomère méthacrylate de dialkylaminoalkyle, ou son sel d'ammonium qua-

ternaire ou son sel d'addition d'acide, (C) est un acrylamide, l est un nombre entier de 0 ou plus, m est un nombre entier de 1 ou plus et n est un nombre entier de 0 ou plus, dans lequel ledit polymère contient un agent de réticulation;

(c) de 0,05% à 5% d'un tensioactif non ionique de rapport hydrolipophile (HLB) élevé, choisi parmi les alcools éthoxylés comportant de 8 à 22 atomes de carbone et de 6 à 30 groupes oxyde d'éthylène; et

(d) de 0,1% à 25% d'un éther alcoxylé de formule:



dans laquelle R est un groupe C<sub>1</sub> à C<sub>18</sub> à chaîne linéaire ou ramifiée, et X est un groupe méthyle ou éthyle, et dans laquelle n a une valeur moyenne de 5 à 20;

la composition étant essentiellement exempte d'éthanol.

2. Composition selon la revendication 1, dans laquelle l'agent de réticulation est choisi parmi le méthylènebisacrylamide, le di(méth)acrylate d'éthylèneglycol, le di(méth)acrylamide, l'acrylate de cyanométhyle, l'acrylate de vinyloxyéthyle, le méthacrylate de vinyloxyéthyle, l'allylpentaérythritol, l'éther diallylique de triméthylolpropane, l'allylsaccharose, le butadiène, l'isoprène, le divinylbenzène, le divinylnaphtalène, l'acrylate d'allyle, et leurs mélanges, de préférence le méthylènebisacrylamide.
3. Composition selon la revendication 2, dans laquelle ledit tensioactif non ionique à HLB élevé est choisi parmi le céth-10, le stéareth-21, et leurs mélanges, et dans laquelle ledit éther alcoxylé est choisi parmi l'éther butylique de PPG-14, l'éther stéarylique de PPG-15, et leurs mélanges.
4. Composition selon la revendication 3, dans laquelle ledit principe pharmaceutique actif est choisi parmi les médicaments anti-acné, les antirides, les principes actifs anti-atrophie de la peau, les anti-inflammatoires non stéroïdiens, les anti-inflammatoires stéroïdiens, les agents bronzants sans soleil, les écrans solaires, les agents de cicatrisation des plaies, les agents de blanchiment ou de décoloration de la peau, les médicaments antihistaminiques, les médicaments antitussifs, les médicaments antipruritiques, les médicaments anticholinergiques, les médicaments anti-émétiques et anti-nauséux, les médicaments anorexiques, les médicaments stimulant le système nerveux central, les médicaments anti-arythmiques, les médicaments β-bloquants, les médicaments cardiotoniques, les médicaments antihypertenseurs, les médicaments diurétiques, les médicaments vasodilatateurs, les médicaments vasoconstricteurs, les médicaments anti-ulcéreux, les médicaments anesthésiques, les médicaments antidépresseurs, les médicaments tranquillisants et sédatifs, les médicaments antipsychotiques, les médicaments antimicrobiens, les médicaments antinéoplasiques, les médicaments antipaludéens, les médicaments myorelaxants, les médicaments antispasmodiques, les médicaments antidiarrhéiques et les médicaments à activité osseuse, et leurs mélanges; de préférence dans laquelle ledit médicament antihistaminique est choisi parmi le maléate de chlorphéniramine, le tannate de chlorphéniramine, le chlorhydrate de triprolidine, l'oxalate de triprolidine, le chlorhydrate de diphenhydramine, l'ascorbate de diphenhydramine, le citrate de diphenhydramine, le succinate de doxylamine, le maléate de pyrilamine, le chlorhydrate de pyrilamine, le tannate de pyrilamine, le tartrate de phénindamine, le chlorhydrate de prométhazine, le chlorhydrate de cyproheptadine, le maléate d'azatadine, le fumarate de clemastine, le maléate de carbinoxamine, le chlorhydrate de carbinoxamine, le chlorhydrate de tripélenamine, le citrate de tripélenamine, le maléate de dexchlorphéniramine, le maléate de bromphéniramine et le chlorhydrate de chlorcyclizine, et leurs mélanges; dans laquelle ledit médicament antitussif est choisi dans le groupe constitué par le bromhydrate de dextrométhorphan, le citrate de carbétapentane, le phosphate de codéine et le chlorhydrate de codéine N-oxyde, et leurs mélanges; dans laquelle ledit médicament anticholinergique est choisi dans le groupe constitué par le bromhydrate de scopolamine, le chlorhydrate de scopolamine, le sulfate d'atropine, le mucate d'atropine, le bromhydrate d'homatropine et le chlorhydrate d'homatropine, et leurs mélanges; dans laquelle ledit médicament anti-émétique ou anti-nauséux est choisi parmi le chlorhydrate de cyclizine, le chlorhydrate de méclizine, le chlorhydrate de chlorpromazine et le maléate de chlorpromazine, et leurs mélanges; dans laquelle ledit médicament anorexique est choisi parmi le chlorhydrate de benzphétamine, le chlorhydrate de phentermine, le chlorhydrate de chlorphentermine et le chlorhydrate de ferfluramine, et leurs mélanges; dans laquelle ledit médicament antimicrobien est choisi parmi les médicaments β-lactames, les médicaments quinolones, la ciprofloxacine, la norfloxacine, la tétracycline, l'érythromycine, l'iséthionat d'hexamidine, l'amikacine, le triclosan, la doxycycline, la capréomycine, la chorhexidine, la chlortétracycline, l'oxytétracycline, la clindamycine,

l'éthambutol, le métronidazole, la pentamidine, la gentamicine, la kanamycine, la linéomycine, la méthacycline, la méthénamine, la minocycline, la néomycine, la nêtilmicine, la paromomycine, la streptomycine, la tobramycine, le miconazole et l'amanfadine, leurs sels pharmaceutiquement acceptables et leurs mélanges; dans laquelle ledit médicament anti-arythmique est choisi parmi le chlorhydrate de propranolol, le chlorhydrate de procainamide, le sulfate de quinidine et le gluconate de quinidine, et leurs mélanges; dans laquelle ledit médicament antihypertenseur est choisi parmi le maléate d'énalapril, le chlorhydrate de clonidine, le chlorhydrate d'hydralazine et le sulfate d'hydralazine, et leurs mélanges; dans laquelle ledit médicament anesthésique ou antipruritique est choisi parmi le chlorhydrate de lidocaïne, le chlorhydrate de bupivacaïne, le chlorhydrate de chlorprocaïne, le chlorhydrate de dibucaïne, le chlorhydrate d'etidocaïne, le chlorhydrate de mepivacaïne, le chlorhydrate de tétracaïne, le chlorhydrate de dyclonine et le chlorhydrate d'hexylcaïne, et leurs mélanges; dans laquelle ledit médicament à activité osseuse est choisi parmi l'acide 6-amino-1-hydroxyhexane-1,1-diphosphonique, l'acide 3-amino-1-hydroxypropane-1,1-diphosphonique, l'acide octahydro-1-pyridine-6,6-diphosphonique, l'acide 2-(2'-pipéridinyl)éthane-1,1-diphosphonique, l'acide 2-(3'-pipéridinyl)éthane-1,1-diphosphonique, l'acide 2-(2'-pipéridinyl)-1-hydroxyéthane-1,1-diphosphonique, l'acide 2-(3'-pipéridinyl)-1-hydroxyéthane-1,1-diphosphonique, l'acide N-(2'-(3'-méthyl)-pipéridinylidène)aminométhanediphosphonique, l'acide N-(2'-(1',3'-diazinylidène))aminométhanediphosphonique et l'acide N-(2-(3-méthylpipéridinylidène))aminométhanediphosphoéthylphosphinique, ou leurs esters et mélanges; dans laquelle ledit anti-inflammatoire non stéroïdien est choisi parmi les dérivés d'acide propionique, les dérivés d'acide acétique, les dérivés d'acide fénamique, les dérivés d'acide biphenylcarboxylique et les oxicames, et leurs mélanges; et dans laquelle ledit anti-inflammatoire non stéroïdien est un dérivé d'acide propionique choisi parmi l'aspirine, l'acétaminophène, l'ibuprofène, le naproxène, le benoxaprofène, le flurbiprofène, le fénopropène, le fenbutène, le kétoprofène, l'indoprofène, le piroprofène, le carprofène, l'oxaprozine, le pranoprofène, le miroprofène, le tiopropofène, le suprofène, l'alminoprofène, l'acide tiaprofénique, le fluprofène et l'acide bucloxyque, et leurs mélanges.

5. Composition selon la revendication 3, dans laquelle ledit principe pharmaceutique actif est un médicament anti-acné choisi parmi l'acide salicylique, le soufre, le résorcinol, la N-acétylcystéine, l'octopirox, l'acide rétinolique et ses dérivés, le peroxyde de benzoyle, l'érythromycine, le zinc, la tétracycline, l'acide azélaïque et ses dérivés, le phénoxyéthanol et le phénoxypropanol, l'acétate d'éthyle, la clindamycine et la méclocycline, les flavinoides, l'acide lactique, l'acide glycolique, l'acide pyruvique, l'urée, le sulfate de scymnol et ses dérivés, le désoxycholate et le cholate, et leurs mélanges, ou un principe actif antirides ou anti-atrophie de la peau choisi parmi l'acide glycolique, l'acide lactique, l'acide salicylique, et leurs mélanges.

6. Composition selon la revendication 3, dans laquelle ledit principe pharmaceutique actif est choisi parmi l'acide salicylique, l'acide glycolique, l'acide lactique, et leurs mélanges.

7. Composition selon la revendication 5, dans laquelle la quantité de (C) dans le polymère cationique est de 50% à 90% en moles; de préférence dans laquelle l, dans le polymère cationique, est nul et le rapport (B):(C) est de 45:55 à 55:45; mieux encore dans laquelle l et n sont tous deux nuls dans le polymère cationique.

8. Composition selon la revendication 7, qui comprend, en outre, de 3% à 5% de glycérine et de 1% à 10% d'un émoulliant choisi parmi les huiles de silicone volatiles, les huiles de silicone non volatiles, les hydrocarbures hautement ramifiés, et leurs mélanges.

9. Composition selon la revendication 3, dans laquelle ledit principe médicamenteux actif est un agent bronzant sans soleil choisi parmi la dihydroxyacétone, les dérivés d'indole, et leurs mélanges; de préférence dans laquelle ladite composition comprend, en outre, un principe actif écran solaire choisi parmi le p-méthoxycinnamate de 2-éthylhexyle, le N,N-diméthyl-p-aminobenzoate de 2-éthylhexyle, l'acide p-aminobenzoïque, l'acide 2-phénylbenzimidazole-5-sulfonique, l'octocrylène, l'oxybenzone, le salicylate d'homomenthyle, le salicylate d'octyle, le 4,4'-méthoxy-t-butylidibenzoylméthane, le 4-isopropylidibenzoylméthane, le 3-benzylidénecamphre, le 3-(4-méthylbenzylidène)camphre, le dioxyde de titane, l'oxyde de zinc, la silice, l'oxyde de fer, et leurs mélanges.

10. Composition pharmaceutique à usage local présentant une pénétration de la peau améliorée, comprenant:


(a) une quantité sans danger et efficace d'un principe pharmaceutique actif,  
 (b) de 0,1% à 10,0% d'un polymère cationique réticulé de masse moléculaire élevée, de formule:  $(A)_l(B)_m(C)_n$ , dans laquelle (A) est un monomère acrylate de dialkylaminoalkyle, ou son sel d'ammonium quaternaire ou son sel d'addition d'acide, (B) est un monomère méthacrylate de dialkylaminoalkyle, ou son sel d'ammonium quaternaire ou son sel d'addition d'acide, (C) est un acrylamide, l est un nombre entier de 0 ou plus, m est un nombre entier de 1 ou plus et n est un nombre entier de 0 ou plus, dans lequel ledit polymère contient un agent de

réticulation;

(c) de 0,05% à 5% d'un tensioactif non ionique de rapport hydrolipophile (HLB) élevé, choisi parmi les alcools éthoxylés comportant de 8 à 22 atomes de carbone et de 6 à 30 groupes oxyde d'éthylène; et

(d) de 0,01% à 5% d'un tensioactif non ionique de HLB faible; et

(e) de 0,1% à 25% d'un éther alcoxylé de formule:



dans laquelle R est un groupe C<sub>1</sub> à C<sub>18</sub> à chaîne linéaire ou ramifiée, et X est un groupe méthyle ou éthyle, et dans laquelle n a une valeur moyenne de 5 à 20;

la composition étant essentiellement exempte d'éthanol.

11. Composition selon la revendication 10, dans laquelle l'agent de réticulation est choisi parmi le méthylènebisacrylamide, le di(méth)acrylate d'éthylèneglycol, le di(méth)acrylamide, l'acrylate de cyanométhyle, l'acrylate de vinyloxyéthyle, le méthacrylate de vinyloxyéthyle, l'allylpentaérythritol, l'éther diallylique de triméthylolpropane, l'allylsaccharose, le butadiène, l'isoprène, le divinylbenzène, le divinylnaphtalène, l'acrylate d'allyle, et leurs mélanges, de préférence dans laquelle ledit agent de réticulation est le méthylènebisacrylamide.

12. Composition selon la revendication 11, dans laquelle ledit tensioactif non ionique à HLB élevé est choisi parmi le céteth-10, le stéareth-21, et leurs mélanges, et dans laquelle ledit éther alcoxylé est choisi parmi l'éther butylique de PPG-14, l'éther stéarylique de PPG-15, et leurs mélanges.

13. Composition selon la revendication 12, dans laquelle ledit tensioactif non ionique à HLB faible est choisi parmi les alcools éthoxylés comportant de 8 à 22 atomes de carbone et de 1 à 5 groupes oxyde d'éthylène; de préférence dans laquelle ledit tensioactif non ionique à HLB faible est choisi parmi le stéareth-2, le céteth-2, le laureth-2, l'oleth-2, et leurs mélanges; et mieux encore dans laquelle ledit tensioactif non ionique à HLB faible est le stéareth-2.

14. Composition selon la revendication 13, dans laquelle ledit principe pharmaceutique actif est choisi parmi les médicaments anti-acné, les antirides, les principes actifs anti-atrophie de la peau, les anti-inflammatoires non stéroïdiens, les anti-inflammatoires stéroïdiens, les agents bronzants sans soleil, les écrans solaires, les agents de cicatrisation des plaies, les agents de blanchiment ou de décoloration de la peau, les médicaments antihistaminiques, les médicaments antitussifs, les médicaments antipruritiques, les médicaments anticholinergiques, les médicaments anti-émétiques et anti-nauséeux, les médicaments anorexiques, les médicaments stimulant le système nerveux central, les médicaments anti-arythmiques, les médicaments β-bloquants, les médicaments cardiotoniques, les médicaments antihypertenseurs, les médicaments diurétiques, les médicaments vasodilatateurs, les médicaments vasoconstricteurs, les médicaments anti-ulcéreux, les médicaments anesthésiques, les médicaments antidépresseurs, les médicaments tranquillisants et sédatifs, les médicaments antipsychotiques, les médicaments antimicrobiens, les médicaments antinéoplasiques, les médicaments antipaludéens, les médicaments myorelaxants, les médicaments antispasmodiques, les médicaments antidiarrhéiques et les médicaments à activité osseuse, et leurs mélanges.

15. Composition selon la revendication 13, dans laquelle ledit principe pharmaceutique actif est un médicament anti-acné choisi parmi l'acide salicylique, le soufre, le résorcinol, la N-acétylcystéine, l'octopirox, l'acide rétinolique et ses dérivés, le peroxyde de benzoyle, l'érythromycine, le zinc, la tétracycline, l'acide azélaïque et ses dérivés, le phénoxyéthanol et le phénoxypropanol, l'acétate d'éthyle, la clindamycine et la méclocycline, les flavinoides, l'acide lactique, l'acide glycolique, l'acide pyruvique, l'urée, le sulfate de scymnol et ses dérivés, le désoxycholate et le cholate, et leurs mélanges, ou un principe actif antirides ou anti-atrophie de la peau choisi parmi l'acide glycolique, l'acide lactique, l'acide salicylique, et leurs mélanges.

16. Composition selon la revendication 13, dans laquelle ledit principe pharmaceutique actif est choisi parmi l'acide salicylique, l'acide lactique, l'acide glycolique, et leurs mélanges.

17. Composition selon la revendication 13, dans laquelle la quantité de (C) dans le polymère cationique est de 50% à

90% en moles; de préférence dans laquelle l, dans le polymère cationique, est nul et le rapport (B):(C) est de 45:55 à 55:45; mieux encore dans laquelle l et n sont tous deux nuls dans le polymère cationique.

- 5 18. Composition selon la revendication 17, qui comprend, en outre, de 3% à 5% de glycérine et de 1% à 10% d'un émollient choisi parmi les huiles de silicone volatiles, les huiles de silicone non volatiles, les hydrocarbures hautement ramifiés, et leurs mélanges.
- 10 19. Composition selon la revendication 13, dans laquelle ledit principe médicamenteux actif est un agent bronzant sans soleil choisi parmi la dihydroxyacétone, les dérivés d'indole, et leurs mélanges; de préférence dans laquelle ladite composition comprend, en outre, un principe actif écran solaire choisi parmi le p-méthoxycinnamate de 2-éthylhexyle, le N,N-diméthyl-p-aminobenzoate de 2-éthylhexyle, l'acide p-aminobenzoïque, l'acide 2-phénylbenzimidazole-5-sulfonique, l'octocrylène, l'oxybenzone, le salicylate d'homomenthyle, le salicylate d'octyle, le 4,4'-méthoxy-t-butyl-dibenzoylméthane, le 4-isopropyl-dibenzoylméthane, le 3-benzylidène-camphre, le 3-(4-méthylbenzylidène)-camphre, le dioxyde de titane, l'oxyde de zinc, la silice, l'oxyde de fer, et leurs mélanges.
- 15

20

25

30

35

40

45

50

55